INDEX VOLUME 53, 1960

Absorbents, effect of on
  plant sugars, 491
  two-spotted spider mite, 491
Abundance
  of bumble bees, 679
  of common carpenter bee, 679
AC-5727, as soil treatment against eye gnat, 1102
Aceratagallia sanguinolenta, 113
Acerophagus notativentris, 757
Acetaldehyde
  as synergist for barthrin in house fly, sprays against, 284
  as synergist in house fly sprays, 300
Acetate of 2,4-dinitro-6-phenyl phenol, toxicity of to oriental fruit moth, 470
Acethion
  metabolic and toxicological relationships of, 25
  rate of detoxification by mosquito of, 779
  resistance of mosquito to, 778
Acethion, toxicity of to
  cockroach, 25
  house fly, 25
  mice, 25
Acethion, dimethyl homolog of
  resistance of mosquito to, 778
Acetic acid
  as synergist for barthrin in sprays against house fly, 284
Acetoacetanilide, effect of on house fly development, 803
Acetone, as attractant to gypsy moth males, 173
Acetophenetidide, effect of on house fly development, 803
Acetophenone, effect of on house fly development, 803
Acetyl phenylhydrazine, effect of on
  fertility of house fly, 804
  house fly development, 803
Acheta
  assimilis, 904
  domesticus, 171
Acid lead arsenate, for control of grape leaf folder, 531
Acrididae, on Ladino clover, 114
Acrobasis vaccinii, 1116
Acrolopphus spp., 294
Acroneuria, 609
Acrylic acid, as synergist in house fly sprays, 300
Acrylonitrile
  effectiveness of against stored corn pests, 238
  fumigation of khapra beetle larvae with, 698
Adalia bipunctata, 659
Adelphocoris
  lineolatus, 823
  rapidus, 527, 823
Aedes
  aegypti, 107, 121, 169, 692, 910, 915, 1037, 1049
  nigromaculis, 876, 1122
  taeniorhynchus, 107
  vexans, 620
Aerosols
  allethrin, against house fly, 956
  Dibrom, against house fly, 956
  studies on fine-particle, coarse-particle, concentrated, 908
Agallia constricta, 114
Agalliinae spp., 823
Age
  effect of on susceptibility of boll weevil, 534
  plant, influence on survival of alfalfa varieties differing in resistance to spotted alfalfa aphid, 142
Ageneotettix deorum, 958
Agriotes
  fuscicollis, 1018
  persimilis, 1018
Agromyza sp., 1133
Agrotis
  fucosa, 1017
  ipsilon, 318
  ypsilon, 527
Ahasverus advena, 536
Alabama argillacea, 104, 242, 293, 747
Aldrin
  against eye gnat, 367
  epoxidation of in soils, 192
  infrared determination of in treated soil, 1129
  long-term effectiveness against eye gnat, 785
  loss of from soil surface, 650
  persistence of in soils, 136
  solubilization of by lipoprotein, 34
Aldrin, effect of on
  boll weevil, 294
  fertility of house fly, 804
  house fly development, 803
  lesser cornstalk borer, 665
  Sacadodes, 294
  seed-corn maggot, 1016
  turnip maggot, 1017
Aldrin, for control of
  boll weevil, 229
  clover root borer, 449
  corn root webworm, 670
  cutworms, 526
  desert grasshopper, 724
  European pine shoot moth, 366
  eye gnats, 362
  fuller rose beetle, 164
  imported fire ant, 190
  Japanese beetle, 320
  lygus bugs, 398
  Nantucket pine moth, 366
  plum curculio, 335, 439
  rice water weevil, 1135
  root maggots, 65
  sugar-beet wireworm, 177
  tarnished plant bug, 690
  woolly apple aphid, 217
Aldrin, residues of
  in soil, 194
  on lima bean, 833
  on radish, 832
  on sweet potato, 833
Aldrin, toxicity of to
  Chrysops, 263
  corn earworm, 22
  house fly, 888
  oriental fruit moth, 470
Aleochara bilineata, 65, 932
Alfalfa
  bee poisoning in, 1010
  effect of hydrogen phosphide on seed germination of, 1
  establishment of parasites of spotted alfalfa aphid in, 1094
  feeding habits of spotted alfalfa aphid on, 714
  influence of plant age on survival of in varieties differing in resistance to spotted alfalfa aphid, 142
  persistence of insecticide residues on, 960
  pollination of by honey bee, 56
  resistance of to spotted alfalfa aphid, 234, 667
  seed damage to by clover seed chalcid, 611
  selection and evaluation of pea aphid-resistant plants, 881
  toxicity of cellulose acetate to, 484
Alfalfa, residues on
  after treatment with phorate, 306
  of malathion, 495
Alfalfa, sampling of
  for clover seed chalcid damage, 611
  plots for insecticide residues, 1078
Alfalfa mosaic, vectors of in clover, 113
Alkoxyphenyl *N*-methylcarbamates, as insecticides, 828
O-Alkyl *S*-2-(ethylthio)-ethyl alkylphosphonothiolate, as systemic insecticide, 127
Allethrin
  aerosols against house fly, 956
  as active ingredient in aerosols, 908
  as spray against house fly, 299
  as synergist for barthrin in sprays against house fly, 284
  effect of on fertility of house fly, 804
Allodermanyssus sanguineus, 476
Allograpta obliqua, 92
Alphitobius diaperinus, 537
Alphitophagus bifasciatus, 537
Altitude, high, exposure to of
  house fly, 247
  oriental rat flea, 247
Amblyomma
  americanum, 953, 1125
  maculatum, 6, 692, 849
American Cyanamid 12008, effect of on survival of serpentine-leaf-miner parasite, 872
Am. Cyanamid 12008, for control of
  cattle grub, 541
  Hessian fly, 501
  serpentine leaf miner, 873
Am. Cyanamid 18133, for control of
  corn earworm, 695
  sand wireworm, 964
Am. Cyanamid 18706
  as soil treatment against eye gnat, 1102
  effect of on plant emergence, 226
  toxicity of to house fly, 741
Am. Cyanamid 18706, effectiveness of against
  cattle grubs, 1126
  lone star tick, 1126
  screw-worms, 1126
  stable fly, 1126
Am. Cyanamid 18706, for control of
  green peach aphid, 1099
  thrips, 227
  two-spotted spider mite, 226
Amethopterin, effect of on
  fertility of house fly, 804
  house fly metabolism, 802
9-Aminoacridine HCl, for control of cattle grubs, 183
9-Amino-6-chloro-2-methoxyacridine, effectiveness of
  against cattle grub, 184
  for control of cattle grubs, 183
Amphimallon majalis, 222
Anagyrus antoninae, 694
Anagyrus diversicornis, 694

INDEX VOLUME 53, 1960

Absorbents, effect of on
  plant sugars, 491
  two-spotted spider mite, 491
Abundance
  of bumble bees, 679
  of common carpenter bee, 679
AC-5727, as soil treatment against eye gnat, 1102
Aceratagallia sanguinolenta, 113
Acerophagus notativentris, 757
Acetaldehyde
  as synergist for barthrin in house fly, sprays against, 284
  as synergist in house fly sprays, 300
Acetate of 2,4-dinitro-6-phenyl phenol, toxicity of to oriental fruit moth, 470
Acethion
  metabolic and toxicological relationships of, 25
  rate of detoxification by mosquito of, 779
  resistance of mosquito to, 778
Acethion, toxicity of to
  cockroach, 25
  house fly, 25
  mice, 25
Acethion, dimethyl homolog of
  resistance of mosquito to, 778
Acetic acid
  as synergist for barthrin in sprays against house fly, 284
Acetoacetanilide, effect of on house fly development, 803
Acetone, as attractant to gypsy moth males, 173
Acetophenetidide, effect of on house fly development, 803
Acetophenone, effect of on house fly development, 803
Acetyl phenylhydrazine, effect of on
  fertility of house fly, 804
  house fly development, 803
Acheta
  assimilis, 904
  domesticus, 171
Acid lead arsenate, for control of grape leaf folder, 531
Acrididae, on Ladino clover, 114
Acrobasis vaccinii, 1116
Acrolopphus spp., 294
Acroneuria, 609
Acrylic acid, as synergist in house fly sprays, 300
Acrylonitrile
  effectiveness of against stored corn pests, 238
  fumigation of khapra beetle larvae with, 698
Adalia bipunctata, 659
Adelphocoris
  lineolatus, 823
  rapidus, 527, 823
Aedes
  aegypti, 107, 121, 169, 692, 910, 915, 1037, 1049
  nigromaculis, 876, 1122
  taeniorhynchus, 107
  vexans, 620
Aerosols
  allethrin, against house fly, 956
  Dibrom, against house fly, 956
  studies on fine-particle, coarse-particle, concentrated, 908
Agallia constricta, 114
Agalliinae spp., 823
Age
  effect of on susceptibility of boll weevil, 534
  plant, influence on survival of alfalfa varieties differing in resistance to spotted alfalfa aphid, 142
Ageneotettix deorum, 958
Agriotes
  fuscicollis, 1018
  persimilis, 1018
Agromyza sp., 1133
Agrotis
  fucosa, 1017
  ipsilon, 318
  ypsilon, 527
Ahasverus advena, 536
Alabama argillacea, 104, 242, 293, 747
Aldrin
  against eye gnat, 367
  epoxidation of in soils, 192
  infrared determination of in treated soil, 1129
  long-term effectiveness against eye gnat, 785
  loss of from soil surface, 650
  persistence of in soils, 136
  solubilization of by lipoprotein, 34
Aldrin, effect of on
  boll weevil, 294
  fertility of house fly, 804
  house fly development, 803
  lesser cornstalk borer, 665
  Sacadodes, 294
  seed-corn maggot, 1016
  turnip maggot, 1017
Aldrin, for control of
  boll weevil, 229
  clover root borer, 449
  corn root webworm, 670
  cutworms, 526
  desert grasshopper, 724
  European pine shoot moth, 366
  eye gnats, 362
  fuller rose beetle, 164
  imported fire ant, 190
  Japanese beetle, 320
  lygus bugs, 398
  Nantucket pine moth, 366
  plum curculio, 335, 439
  rice water weevil, 1135
  root maggots, 65
  sugar-beet wireworm, 177
  tarnished plant bug, 690
  woolly apple aphid, 217
Aldrin, residues of
  in soil, 194
  on lima bean, 833
  on radish, 832
  on sweet potato, 833
Aldrin, toxicity of to
  Chrysops, 263
  corn earworm, 22
  house fly, 888
  oriental fruit moth, 470
Aleochara bilineata, 65, 932
Alfalfa
  bee poisoning in, 1010
  effect of hydrogen phosphide on seed germination of, 1
  establishment of parasites of spotted alfalfa aphid in, 1094
  feeding habits of spotted alfalfa aphid on, 714
  influence of plant age on survival of in varieties differing in resistance to spotted alfalfa aphid, 142
  persistence of insecticide residues on, 960
  pollination of by honey bee, 56
  resistance of to spotted alfalfa aphid, 234, 667
  seed damage to by clover seed chalcid, 611
  selection and evaluation of pea aphid-resistant plants, 881
  toxicity of cellulose acetate to, 484
Alfalfa, residues on
  after treatment with phorate, 306
  of malathion, 495
Alfalfa, sampling of
  for clover seed chalcid damage, 611
  plots for insecticide residues, 1078
Alfalfa mosaic, vectors of in clover, 113
Alkoxyphenyl *N*-methylcarbamates, as insecticides, 828
O-Alkyl *S*-2-(ethylthio)-ethyl alkylphosphonothiolate, as systemic insecticide, 127
Allethrin
  aerosols against house fly, 956
  as active ingredient in aerosols, 908
  as spray against house fly, 299
  as synergist for barthrin in sprays against house fly, 284
  effect of on fertility of house fly, 804
Allodermanyssus sanguineus, 476
Allograpta obliqua, 92
Alphitobius diaperinus, 537
Alphitophagus bifasciatus, 537
Altitude, high, exposure to of
  house fly, 247
  oriental rat flea, 247
Amblyomma
  americanum, 953, 1125
  maculatum, 6, 692, 849
American Cyanamid 12008, effect of on survival of serpentine-leaf-miner parasite, 872
Am. Cyanamid 12008, for control of
  cattle grub, 541
  Hessian fly, 501
  serpentine leaf miner, 873
Am. Cyanamid 18133, for control of
  corn earworm, 695
  sand wireworm, 964
Am. Cyanamid 18706
  as soil treatment against eye gnat, 1102
  effect of on plant emergence, 226
  toxicity of to house fly, 741
Am. Cyanamid 18706, effectiveness of against
  cattle grubs, 1126
  lone star tick, 1126
  screw-worms, 1126
  stable fly, 1126
Am. Cyanamid 18706, for control of
  green peach aphid, 1099
  thrips, 227
  two-spotted spider mite, 226
Amethopterin, effect of on
  fertility of house fly, 804
  house fly metabolism, 802
9-Aminoacridine HCl, for control of cattle grubs, 183
9-Amino-6-chloro-2-methoxyacridine, effectiveness of
  against cattle grub, 184
  for control of cattle grubs, 183
Amphimallon majalis, 222
Anagyrus antoninae, 694
Anagyrus diversicornis, 694

Analog- Synergism, of several carbamate insecticies, 1004
Anarsia lineatella, 680
Anasa tristis, 389
Anastrepha ludens, 323
Anchytarsus, 609
p-Anisic acid, as synergist for barthrin in sprays against house fly, 284
Anopheles
albimanus, 634
aztecus, 634
quadrimaculatus, 101, 121, 634, 915
Ant, carpenter, as carrier of oak wilt, 578
Ants, malathion for reducing dried-fruit packing house infestations of, 970
Anthocoridae, as predators of spotted alfalfa aphid, 92
Anthonomus grandis, 41, 52, 105, 228, 291, 311, 416, 534, 551, 681, 688, 711, 747, 844, 849, 1051
Anthremus verbasci, 537
Anticarsia gemmatalis, 131
Anticide, in baits for control of imported fire ant, 190
Antonina
graminis, 694
purpurea, 694
Anuraphis bakeri, 113, 1012
Anyphaena celer, 811
Anyphaenidae, in apple orchards, 811
Apanteles
autographae, 567
bedelliae, 1119
congregatus, 976
egena, 567
medicaginis, 71
Aphelinus semiflavus, 655, 1094
Aphidencyrtus aphidivorus, 151
Aphidius
alius, 151
confusus, 151
knowltoni, 659
nigripes, 151
Aphis
fabae, 769, 1024
gossypii, 103, 294, 655, 747
medicaginis, 115
ponii, 685
spiraecola, 258
Aphytis, 970, 1033
diaspidis, 757
lingnanensis, 971
mytilaspidis, 757
Apis mellifera, 56, 95, 266, 546, 564, 679, 696, 782, 993, 1010, 1137
Apple
effect of apple seed chalcid on, 166
effect of soil insecticides on, 487
honey bee count on, 565
influence of spray programs on, 717
measuring surface area of for expression of pesticide residues on, 462
soil injection of systemic acaricides for control of mites on, 144
Aqua Nuchar A, effect of on plant sugars and resulting effect on mite nutrition, 491
Arachnophaga frontalis, 305
Aramite, effect of on Pacific spider mite, 626
Aramite, for control of
cotton mite, 203
lygus bugs, 398
mites, 735
myocoptic mange mite, 168
spider mite complex, 522
Araniella diplicata, 812
Arason, effect of on germination of wheat, 575
Argiopidae, in apple orchards, 811
Argyrotaenia velutinana, 695
Arrestant, as standard term for insect reaction to chemicals, 134
Arrhenodes minutus, 510
Arsenicals, for control of desert grasshopper, 724
Asaphes californicus, 151
Aspidiotus perniciosus, 1085
Assay, insecticide, apparatus for, 483
Asterolecanius sp., 406
Atta spp., 294
Attagenus, 537
piceus, 966, 974
Attractant, as standard term for insect reaction to chemicals, 134
Aulocara elliotti, 640
Aureomycin, effect of on house fly larvae, 776
Autophisia egena, 566
2,4,6-tris (1-Aziridinyl)-*s*-triazine, effect of on
fertility of house fly, 804
house fly development, 803

bis (1-Aziridinyl) morphelino-phosphine sulfide, effect of on
fertility of house fly, 804
house fly development, 803
Bacillus thuringiensis, against
cotton leafworm, 747
European corn borer, 624
mint looper, 530
Bacillus thuringiensis, for control of
bean leaf skeletonizer, 566
European pine shoot moth, 352
grape leaf folder, 531
house fly, 774
Bacillus thuringiensis, toxicity of to corn earworm, 22
Baetis, 609
Bait, Dimetilan, for house fly control, 898
Bait, insecticide, for systemic control of fleas on rats, 169
Barley, effect of hydrogen phosphide on germination of, 1
Barthrin, for control of
house fly, 282
poultry lice and mites, 161
Barthrin, toxicity of to
chicken mite, 12
house fly, 741
northern fowl mite, 12
Bayer 16948, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Bayer 16948, toxicity of to
house fly, 6
white rat, 6
Bayer, 18510, toxicity of to
house fly, 6
white rat, 6
Bayer 18779, toxicity of to
house fly, 6
white rat, 6
Bayer 21/199 (*see Co-Ral*)
Bayer 22408
as soil treatment against eye gnat, 1102
biological degradation of, 848
Bayer 22408, for control of
European pine shoot moth, 349
grape leaf folder, 531
lone star tick, 953
screw-worm, 1058
Bayer 22684, effectiveness against
bed bug, 6
Gulf Coast tick, 6
house fly, 6
white rat, 6
Bayer 22893, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Bayer 22893, toxicity of to
house fly, 6
white rat, 6
Bayer 23129
effectiveness of against cattle grubs, 1126
for control of thrips, 225
Bayer 23453, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Bayer 23453, toxicity of to
house fly, 6
white rat, 6
Bayer 24498, effectiveness of against
bed bug, 6
cattle grubs, 1126
Gulf Coast tick, 6
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Bayer 24498, toxicity of to
house fly, 6
white rat, 6
Bayer 24869, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Bayer 24869, toxicity of to
house fly, 6
oriental fruit moth, 470
white rat, 6
Bayer 24882, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Bayer 24882, toxicity of to
house fly, 6
white rat, 6
Bayer 25141, against
boll weevil, 747
bollworm, 711, 747
Bayer 25141
against desert spider mite, 748
as soil treatment against eye gnat, 1102
effect of on plant emergence, 226
toxicity of to oriental fruit moth, 470
Bayer 25141, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Bayer 25141, for control o
boll weevil, 1052
cotton aphid, 747
cotton fleahopper, 1052
green peach aphid, 404
spider mite, 1053
thrips, 227, 1051
two-spotted spider mite, 226
Bayer 25142, effectiveness of agains
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Bayer 25198, effectiveness of against
bed bug, 6
cattle grubs, 1126
Gulf Coast tick, 6
lone star tick, 1126
stable fly, 1126
screw-worms, 1126
Bayer 25198, toxicity of to
house fly, 6
white rat, 6
Bayer 29493
as soil treatment against eye gnat, 1102
effect of on plant emergence, 226

Bayer 29493, against
boll weevil, 711
bollworm, 712
Bayer 29493, effectiveness of against
bed bug, 6
cattle grubs, 1126
Gulf Coast tick, 6
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Bayer 29493, for control of
European pine shoot moth, 352
green peach aphid, 404
house fly, 539
two-spotted spider mite, 226
thrips, 227
Bayer 29493, toxicity of to
house fly, 6, 741
white rat, 6
Bayer L 13/59 (*see Dipterex and Dylox*)
Bean, Black Valentine
effect of absorbents on, 491
effect of soil insecticides on, 491
Bean, broad, toxicity of cellulose acetate to, 484
Bean, green, residues of nicotine on, 64
Bean, kidney, seed-corn maggot damage to seed of, 1016
Bean, lima, systemic insecticidal properties of carbamates on, 1021
Bean yellow mosaic, vectors of in clover, 113
Bedellia somnulentella, 1119
Bee beds, artificial, for propagation of alkali bee, 1025
Bee poisoning, vs clover aphid control in red clover, 1012
Bees, wild, abundance of, 679
Benzene, as attractant to gypsy moth males, 173
Benzene, as synergist for barthrin in sprays against house fly, 284
Benzeneboronic acid, effect of on fertility of house fly, 804
Benzenesulfonate ester, synergistic effect upon mortality of
DDT-resistant house fly from Sevin, 232
parathion-resistant house fly from Sevin, 232
Benzoic acid, as synergist for barthrin in sprays against house fly, 284
beta-Benzyloxyethyl acrylate, effect of on house fly development, 803
Berries, wild, as forage for bees, 679
BHC
persistence of residues on alfalfa, 960
residues of in mint oil, 531
translocation from treated soil, 110
BHC, effect of on
apple, 487
boll weevil, 294
germination of red clover, 110
growth of red clover, 110
mite nutrition, 487
Sacadodes, 294
seed-corn maggot, 1016
BHC, for control of
boll weevil, 229
clover root borer, 449
European pine shoot moth, 353
Japanese beetle, 320
woolly apple aphid, 217
BHC, toxicity of to
boll weevil, 534
loblolly pine seedlings, 175
vedalia beetle, 1056
Biology of
apple seed chalcid, 166
black cherry aphid, 659
blackheaded fireworm of cranberry, 1097
boll weevil, 41
brown wheat mite, 704
convergent lady beetle when fed diet of spotted alfalfa aphid, 257
European pine shoot moth, 349
Hyperodes humilis, 252
morning-glory leaf miner, 1119
little house fly, 999
oak timberworm, 510
peach twig borer, 680
periodical cicada, 961
pine webworm, 587
predators of spotted alfalfa aphid, 89
Psorophora cyanescens, 619
San Jose scale, 757
Solierella sp. 958
tarnished plant bug, 1063
three-lined potato beetle, 481
white-pine shoot borer, 249
Biological control, of cottony-cushion scale, 1055
Bionomics, of *Psorophora cyanescens*, 619
Black Leaf 40 (*see Nicotine*)
Blapstinus sp, 970
Blatta orientalis, 170, 904
Blattella germanica, 200, 806, 849, 904, 1004, 1038, 1138
Blissus leucopterus, 823
Bombus, 546, 680, 1010
borealis, 548
rufocinctus, 548
terricola, 548
vagans, 548
Book Review
Biological and Chemical Control of Plant and Animal Pests, 1141
Chemical and Natural Control of Pests, 1140
Cholinesterases: A Histochemical Contribution to the Solution of Some Functional Problems, 569
Diseases and Pests of Ornamental Plants, 977
Grasslands, 179
Mosquitoes of Medical Importance, 330
A Synoptic Catalog of the Mosquitoes of the World, 330
A Review of the Biological Control of Insects and Weeds in Australia and Australian New Guinea, 1140
Subtropical Fruit Pests, 179
Ticks—A Monograph of the Ixodoidea, 1140
Tyroglyphoidea (Acari). Fauna of USSR. Arachnoidea, 554
Bracon hebetor, 537
Bradysia fenestralia, 558
Breeding, amount of by *Drosophila* in tomato refuse, 8
Brevicoryne brassicae, 47, 389
5-Bromouracil, effect of on house fly development, 803
Bruchophagus gibbus, 611
Bruchus brachialis, 537, 555
Bryobia arborea, 522
Bucculatrix thurberiella, 76, 127
2,3,4,5—Bis (Δ^2-butenylene) tetrahydrofurfural, screening of as cockroach repellent, 806
Butonate, effectiveness of against
screw-worms, 1126
cattle grubs, 1126
lone star tick, 1126
stable fly, 1126
m-n-Butoxy-*N*-methylcarbamate, toxicity of to salt-marsh caterpillar, 1022
m-sec. Butoxy-*N*-methylcarbamate, toxicity of to salt-marsh caterpillar, 1022
Butoxy polypropylene glycol, repellency of to
horn fly, 435
stable fly, 434
bis (tri-*n*-Butyltin) oxide, against house fly, 447
n-Butyltin-tri (2-ethylhexanoate), against house fly, 447
m-tert-Butyl-*N*-methylcarbamate, toxicity of to salt-marsh caterpillar, 1022
m-t-Butylphenyl *N*-methylcarbamate, as soil treatment against eye gnat, 1104
tert-Butylsulfenyl dimethyl dithiocarbamate, screening of as cockroach repellent, 806

C-140
for bee poisoning, 1013
for control of clover aphid, 1013
C-3975, toxicity of to corn earworm, 22
C-4018, toxicity of to corn earworm, 22
C-6812, toxicity of to corn earworm, 22
C-8305, toxicity of to corn earworm, 22
C-25141, toxicity of to corn earworm, 22
Cabbage, uptake of phorate by, 47
Cabinet, temperature, for bio-ecological studies, 1030
Calcium arsenate
tolerance of boll weevils to, 844
toxicity of to vedalia beetle, 1056
Calcium arsenate, effect of on
boll weevil, 294
Sacadodes, 294
Calcium arsenate, for control of
boll weevil, 228
potato tuberworm, 868
Calcium hydroxide, effect of as absorbent on
plant sugars, 491
mite nutrition, 491
Caleida decora, 567
Calliephialtes grapholithae, 305
Callitroga hominivorax, 1058, 1110, 1125, 1133
Calomycterus setarius, 865
Campodeidae, 823
Campoplex sp., 585
Captan, effect of on germination of wheat, 575
Captan, for control of
European red mite, 892
mites, 735
Captan, influence of soil type and moisture on wheat treated with, 773
Carabidae, abundance of in dieldrin-treated areas, 823
Carbamates, N-Alkyl
esterase reaction mechanism of, 205
stability of, 205
toxicity of to house fly, 205
toxicity of to mouse, 205
Carbamates, N,N-dialkyl
esterase reaction mechanism of, 205
stability of, 205
toxicity of to house fly, 205
toxicity of to mouse, 205
Carbon bisulphide, in control of stored-grain insects, 536
Carbon disulfide
distribution and sorption of applied to wheat by recirculation, 259
for control of rice weevil, 288
Carbon tetrachloride
distribution and sorption of applied to wheat by recirculation, 259
effectiveness of against stored corn pests, 238
for control of rice weevils, 288
for control of stored-grain insects, 536

fumigation effect of on confused flour beetle, 1071
fumigation effect of on rice weevil, 1071
Carpocapsa pomonella, 685, 695, 1083
Carpophilus, 970
dimidiatus, 389
hemipterus, 537
lugubris, 174
mutilatus, 537
sayi, 537
Cats, effect of dieldrin on, 826
Cattle, metabolism of P^{32}-Delnav in, 672
Celama sorghiella, 749
Cellulose acetate, toxicity of to leguminous plants, 484
Cephalonomia, 970
tarsalis, 537
waterstoni, 537
Cephus cinctus, 244
Ceramica picta, 527
Ceraticelus fissiceps, 812
Ceratocystus
fagearum, 578
ulmi, 1034
Cereals
effect of hydrogen phosphide on seed germination of, 1
X-ray microscope for study of internal infesting insects of, 678
Chaetocnema
confinis, 822
denticulata, 132, 527, 822
pulicaria, 527, 822
Chaitophorus negundinis, 258
Charcoal, effect of as absorbents on plant sugars, 491
Charips, spp., 151
Chemagro 102, effect of on plant emergence, 226
Chemagro 102, for control of
thrips, 225
two-spotted spider mite, 226
Chemagro 103, effect of on plant emergence, 226
Chemagro 103, for control of
thrips, 225
two-spotted spider mite, 226
Chemagro 104, effect of on plant emergence, 226
Chemagro 104, for control of
thrips, 225
two-spotted spider mite, 226
Chermes abietis, 979
Chick, effect of dipterex-treated water on, 1067
Chipman R-6199
effectiveness of soil injection of against European red mite, 145
effectiveness of soil injection of against two-spotted spider mite, 145
for control of introduced pine sawfly, 648
penetration of cotton leaves by, 106
residues of in cotton leaves, 106
toxicity of to oriental fruit moth, 469
Chipman R-6199, residual effect of on
boll weevil, 105
bollworm, 104
cabbage looper, 104
cotton aphid, 105
cotton leafworm, 104
mealy bugs, 105
pink bollworm, 105
salt-marsh caterpillar, 104
soft scale, 104
thrips, 105
two-spotted spider mite, 105
whitefly, 104
Chironomidae, survival of after DDT spraying, 609
Chlamydatus associatus, 823
Chlorbenside
against two-spotted spider mite, 841
for control of a spider mite complex, 522
Chlordane
against eye gnat, 367
against raisin moth, 975
development of resistance to in cockroach, 1138
duration of toxicity of to fuller rose beetle, 165
in mint insect control, 529
Chlordane, effect of soil treatments of on flavor of
carrots, 285
parsnips, 285
turnips, 285
Chlordane, for control of
clover root borer, 449
corn earworm, 398
corn root webworm, 670
cutworms, 526
desert grasshopper, 724
European pine shoot moth, 366
fuller rose beetle, 164
imported fire ant, 190, 647
Japanese beetle, 320
lygus bugs, 398
mint insects, 527
Nantucket pine moth, 366
plum curculio, 439
sugar-beet wireworm, 177
woolly apple aphid, 217
Chlordane, residues of in on
mint oil, 531
vegetable crops, 832
Chlordane, toxicity of to *Chrysops*, 263
β-chlordane, toxicity of to house fly, 889
Chlorobenzilate, duration of toxicity of to fuller rose beetle, 165
Chlorochroa
ligata, 689
sayi, 75, 689
2,5-bis [N,N-bis (2-Chloroethyl)-aminomethyl] hydroquinone, effect of on house fly development, 803
Chloroferron
anticholinesterase activity of, 1039
toxicity of, 1039
Chloroplast, number in
Hessian-fly infested wheat leaves, 560
maleic hydrazide-treated wheat leaves, 560
Chlorthion
duration of toxicity of to fuller rose beetle, 165
effectiveness of on salt-marsh caterpillar, 78
for control of plum curculio, 335
residues of in mint oil, 529
resistance of mosquito to, 777
Chlorthion, toxicity of to
Chrysops, 263
corn earworm, 22
oriental fruit moth, 469
vedalia beetle, 1056
Cholinesterase
activity of in blood from cattle fed phorate-treated hay, 309
blood, effect of organic phosphorous insecticides on, 603
erythrocyte and plasma levels of, 593
erythrocyte, following administration of systemic insecticides, 185
Choristoneura
fumiferana, 631, 684
pinus, 587
Chorizagrotis auxiliaris, 319
Chrysanthemumate, toxicity of to house fly, 742
Chrysanthemumic acid, as synergist for barthrin in sprays against house fly, 284
Chrysocharis
ainsliei, 682
parksi, 682
Chrysopa, 68, 152, 661
oculata, 92, 1088
plorabunda, 92
Chrysopidae, as predators of spotted alfalfa aphid, 92
Chrysops flavida v. *reichteri*, 262
Cimex lectularius, 6, 692, 849, 904
Circulifer tenellus, 162
Clover
as forage for bees, 679
bee poisoning vs clover aphid control on, 1012
effect of hydrogen phosphide on seed germination of, 1
effect of root-feeding weevils on longevity of, 865
honey bee count on, 565
honey bee counts in relation to blossoming of, 548
incidence of root-feeding weevils on, 865
pollination of, 546
residues of dieldrin on, 827
resistance to pea aphid of, 863
vectors of mosaic viruses on, 113
Coccinellidae
abundance of in dieldrin-treated area, 823
as predators of spotted alfalfa aphid, 92
Coccinellids, toxicity to of Dylox, 68
Coccophagoides kuwanae, 757
Coccus hesperidum, 104
Coccygomimus inflatus, 305
Coefficients, of co-toxicity of insecticides tested on house fly, 890
Colaspis flavida, 132
Colchicine, effect of on screw-worms, 1133
Coleomegilla
maculata, 1088
m. lengi, 91
Coleoptera, attracted to black light, 173
Colias philodice eurytheme, 73, 319
Collembola, effect of insecticides on, 358
Collops vittatus, 70
Colonization, of mosquitoes, 101
Complementary activity, of acaricides, 967
Compound 7769, for control of boll weevil, 229
Conotrachelus nenuphar, 335, 439
Conozoa
carinata, 722
sulcifrons sulcifrons, 722
Contarinia sorghicola, 749
Copidosoma truncatellum, 566
Copper compounds, and zineb for control of citrus rust mites, 395
Copper-sulfur-lime, influence of on growth of apple, 717
Co-Ral (Bayer 21/199)
against stable fly, 303
animal systemic activity of in rabbits, 1037
anticholinesterase activity of, 1039
as soil treatment against eye gnat, 1102
determination of in animal tissues, 263
histopathological effects of on gullet tissues of
cattle, 280
metabolism of, 1037
resistance of mosquito to, 777
stability of, 1037
Co-Ral, for control of
cattle grubs, 125, 183, 541, 683, 729
lone star tick, 953

northern fowl mite, 4
poultry lice and mites, 160
screw-worm, 1058
Co-Ral, toxicity of to
bed bug, 1039
Chrysops, 263
Gulf Coast tick, 1037
horses, 602
house fly, 429, 1039
mosquitoes, 1039
rats, 1039
Co-Ral, *O*-analog of
anticholinesterase activity of, 1039
toxicity of, 1039
Co-Ral, radioactive, fractionation of, 1040
Cords, Dimetilan-impregnated, for house fly control, 898
Cordulegaster, 609
Coriarchne sp.
Corixidae, attracted to black light, 173
Corizus lateralis, 527
Corn
heptachlor residues on stover of, 19
in dieldrin-treated fields, 584
loss in yield of due to corn leaf aphid, 924
Corn, effect on of
European corn borer, 517
hydrogen phosphide, on germination of, 1
Sevin, 388
sugarcane borer on yield of, 935
Corn, field, resistance of to European corn borer, 918
Corn, high-amylose, resistance of to Angoumois grain moth, 573
Corn, stored, fumigation of, 107
Corn oil, effect of on house fly, 325
Corn silk, effect of DDT on egg-predator complex of corn earworm in, 1088
Corticaria sp., 537
Cotton
effect of dieldrin on, 826
flight distance of boll weevil on, 682
insects of in Nicaragua, 291
nectariless, effect of on lepidopterus populations, 242
phytotoxicity study on, 203
sampling techniques for lygus and stink bugs on, 689
tolerance of boll weevils on, 844
toxicity of DDVP to, 593
Crambus caliginosellus, 670
Cremnops vulgaris, 585
Cryolite, for control of
cranberry fruitworm, 1116
fuller rose beetle, 164
Cryptochaetum iceryae, 1055
Cryptophagus sp., 537
Cryptoserphus abruptus, 537
Cryptotermes brevis, 503
Ctenicera puncticollis, 1018
Ctenocephalides felis, 904
Cue-lure, as new common name of insecticide, 677
Culex
fatigans, 915
peus, 876
pipiens quinquefasciatus, 876, 1123
quinquefasciatus, 650, 828, 938
tarsalis, 876, 904, 1122
Culicoides variipennis sonorensis, 731
Culm solidness, effect of on survival of wheat stem sawfly, 244
Curly top, effect of on yield and quality in sugar beet, 162
Cutworm, climbing, on mint, 527
Cyclocephala, 822
borealis, 860
munda, 91
polita, 659
Cycloneda munda, 91
Cylindrocopturus eatoni, 1044
Cyrtepistomus castaneus, 132
Cyrtolobus
arcuatus, 173
auroreus, 173
discoidalis, 173
fuliginosis, 173
fuscipennis, 173
helena, 173
maculifrontis, 173
pallidifrontis, 173
querci, 173
tuberosus, 173
vau, 173

Daikon, damage to by turnip maggot, 1017
Dalopius sp., 527
Dandelion, as forage for bees, 679
DDD, reduction of on grapes, 273
DDE
for control of boll weevil, 551
production of by mosquito larvae, 1049
solubilization of by lipoprotein, 34
DDT
action of at an insect synapse, 170
action of on mosquitoes, 107
deposits on grapes as affected by growth and weathering, 269
duration of toxicity to fuller rose beetle, 165
excretion of in milk of cows fed on DDT-treated pastures, 411
in aerosols, 909
influence on growth of apple, 717
interference in biological control of cottony-cushion scale, 1055
long-term effectiveness against eye gnat, 785
loss of from soil surface, 650
persistence of in soils, 136
solubilization of by lipoprotein, 34
DDT, against
Anopheline mosquitoes, 634
boll weevil, 711, 747
bollworm, 711, 747
cotton leafworm, 749
eye gnat, 367
European corn borer, 624
mosquitoes, 121, 1049
raisin moth, 975
spruce budworm, 631
stable fly, 303
DDT, effect of on
apple, 487
aquatic insects, 608
beans, Black Valentine, 491
boll weevil, 294
Clark soybeans, 491
egg-predator complex of corn earworm, 1088
mite nutrition, 487
Sacadodes, 294
soil microarthropods, 357
transamination in cockroach, 167
two-spotted spider mite, 491
yield of soybeans, 133
DDT, effectivenss of on
boll weevil, 53
cotton fleahopper, 53
desert spider mite, 54
salt-marsh caterpillar, 78
DDT, for control of
apple aphid, 718
apple seed chalcid, 166
beet armyworm, 618
bollworm, 104
boll weevil, 229, 1051
cabbage looper, 104
codling moth, 685
corn earworm, 397, 695, 819
cotton fleahopper, 1051
cotton leafworm, 104
cotton mite, 204
cranberry fruitworm, 1118
Drosophila, 1109
European corn borer, 590
European pine shoot moth, 349, 365
European red mite, 892
eye gnats, 362
face fly, 450
grape berry moth, 269, 857
green peach aphid, 403
Hyperodes humilis, 251
imported fire ant, 190, 647
insect pests on tomatoes, 1128
lesser cornstalk borer, 664
lygus bugs, 74, 397, 514, 767
maple leaf cutter, 957
mint insects, 527
mint looper, 527
mites, 735
morning-glory leaf miner, 1119
Nantucket pine moth, 365
pink bollworm, 105
potato leafhopper, 718
potato tuberworm, 868
spider mites, 1053
sugar-beet wireworm, 177
three-lined potato beetle, 480
thrips, 227, 1052
DDT, residual effect of on
salt-marsh caterpillar, 104
DDT, residues of in/on
alfalfa-brome grass pasture, 412
milk, 1019
mint oil, 527
peppers, 622
vegetable crops, 832
DDT, resistance to of
codling moth, 1083
house fly, 313
mosquito, 777, 1128
DDT, toxicity of to
boll weevil, 551
cabbage looper, 991
Chrysopa spp., 69
Chrysops, 263
convergent lady beetle, 69
corn earworm, 22
Geocoris spp., 69
house fly, 38, 709
Nabis ferus, 69
oriental fruit moth, 469
Orius spp., 69
syrphids, 69
Trioxys utilis, 69
vedalia beetle, 1056
DDT, aliphatic deuterium analog of degradation of when applied to house fly, 40
synethsis of, 35
toxicity of to house fly, 35
p,p'-CH_3DDT, solubilization of by lipoprotein, 34
p,p'-Br DDT-solubilization of by lipoprotein, 34
p,p'-F DDT-solubilization of by lipoprotein, 34
p,p'-I DDT, solubilization of by lipoprotein, 34
DDT-dehydrochlorinase III, effect of enzyme concentration on activity of, 33
DDT irritation threshold, relation of to knockdown of Anopheline moquitoes, 634
DDT-oil, for control of sap beetles, 175
DDT resistance, effect of on development

of malathion resistance in house fly, 320
DDT-Risella, susceptibility to of mosquitoes, 103
DDVP
conversion of Dipterex to, 1069
for control of face fly, 450
resistance of mosquito to, 777
DDVP, toxicity of to
cows, 593
horses, 593, 602
house fly, 599, 741, 1069
white rats, 593
Deet, added common name of insecticide, 677
Delnav, as soil treatment against eye gnat, 1102
Delnav, effect of on survival of serpentine-leaf-miner parasite, 872
Delnav, for control of
apple seed chalcid, 166
bean leaf skeletonizer, 567
corn earworm, 401
cranberry fruitworm, 1118
European pine shoot moth, 352, 365
imported fire ant, 190
lone star tick, 953
lygus bugs, 401
Nantucket pine moth, 365
Oligonychus pratensis, 1061
poultry lice and mites, 161
serpentine-leaf-miner, 872
spider mites, 726
Delnav, toxicity of to
corn earworm, 22
house fly, 741
Oriental fruit moth, 470
Delnav, resistance of mosquito to, 777
Delphacinae spp., 823
Delphene, repellency of to
horn fly, 435
stable fly, 434
Demeton
against desert spider mite, 748
effectiveness of soil injection of against European red mite and two-spotted spider mite, 145
for bee poisoning, 1013
influence of on apple growth, 717
resistance of citrus red mites to and cross-tolerance to Kethane and Tedion, 384
resistance of mosquito to, 777
Demeton, effect of on
beetle, 1035
Ladino clover, 1134
plant emergence, 226
plants, 838
survival of serpentine-leaf-miner parasite, 872
Demeton, for control of
apple seed chalcid, 166
clover aphid, 1013
corn earworm, 400
cotton aphid, 747
greenbug, 278
green peach aphid, 727, 835, 1100
Hyperodes humilis, 254
introduced pine sawfly, 648
leafhopper, 835
lygus bugs, 398
Oligonychus pratensis, 1061
potato leafhopper, 718
spider mites, 725
thrips, 225
tobacco flea beetle, 835
two-spotted spider mite, 227
woolly apple aphid, 217
Demeton, toxicity of to
Chrysops, 263
house fly, 741
oriental fruit moth, 470
Dendroctonus
engelmanni, 459
monticolae, 461
Dermanyssus gallinae, 12, 328, 476, 904
Dermestes lardarius, 537
Dermestids, malathion for reducing dried-fruit packing house infestations of, 970
Derocerus reticulatum, 113
Derostemes
arizonensis, 682
variipes, 872
Desmia funeralis, 531
Deterrent, as standard term for insec reaction to chemicals, 134
Development
of sorghum, effect of phorate applied to seed on 154
Diabrotica
longicornis, 822
undecimpunctata howardi, 132, 822
Dialictus
admirandus, 680
inconspicuus, 680
nymphaearum, 680
viridissimus, 680
Diapause, influencing factors in pink bollworm, 793
Diaphania nitidalis, 482
Diatraea saccharalis, 935
Diazinon
duration of toxicity to fuller rose beetle, 165
effectiveness of on salt-marsh caterpillar, 78
resistance of house fly to, 313
Diazinon, for control of
bean leaf skeletonizer, 568
boll weevil, 229
cockroach, 200
codling moth, 1083
corn earworm, 399
cranberry fruitworm, 1116
Drosophila, 1107
European red mite, 892
face fly, 450
grape leaf folder, 533
grape mealybug, 354
green peach aphid, 404
house fly, 315, 539, 898
Hyperodes humilis, 254
imported fire ant, 190
little house fly, 999
lygus bugs, 398, 514
morning-glory leaf miner, 1119
plum curculio, 335
San Jose scale, 1085
sap beetles, 175
serpentine-leaf-miner, 872
spider mite complex, 522
Diazinon, toxicity of to
Chrysops, 263
cockroach, 25
corn earworm, 22
house fly, 25, 429
oriental fruit moth, 469
serpentine-leaf-miner parasite, 872
vedalia beetle, 1056
Dibrachys, 305
cavus, 305
Dibrom
aerosols against house fly, 956
against European corn borer, 625
effect of on survival of serpentine-leaf-miner parasite, 872
for bee poisoning, 1013
Dibrom, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Dibrom, for control of
clover aphid, 1013
cranberry fruitworm, 1118
Drosophila, 1107
green peach aphid, 404
Mexican bean beetle, 819
parasitic mites, 476
tarnished plant bug, 690
Dibrom, toxicity of to
corn earworm, 22
house fly, 6, 741
white rat, 6
Di-*n*-butyl succinate, screening of as cockroach repellent, 806
Di-*n*-butyltin dichloride, against house fly, 447
Di-*n*-butyltin-di(2-ethylhexanoate), against house fly, 447
Di-*n*-butyltin dilaurate
effectiveness of against cattle grubs, 184
for control of cattle grubs, 183
Dicalcium arsenate, effect of on boll weevil, 845
Dicapthon, against
boll weevil, 711
bollworm, 711
eye gnat, 1102
Dicapthon, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
Dicapthon, for control of
boll weevil, 229
lone star tick, 953
poultry lice and mites, 160
Diacapthon, toxicity of to
corn earworm, 22
house fly, 741
northern fowl mite, 12
2,2″-Dichloro-*N*-methyldiethyl-amine hydrochloride, effect of on house fly development, 803
2-2-Dichlorovinyl Dimethyl Phosphate, toxicology of, 593
Dicranota, 609
Dieldrin
against eye gnat, 367
against mosquitoes, 124
duration of toxicity of to fuller rose beetle, 165
excretion of in milk of cows fed on dieldrin-treated pastures, 411
formed from aldrin in soil, 192
for suppression of Japanese beetles, 582
increase of European corn borer following application of, 582
infrared determination of in treated soil, 1129
in protection against bee poisoning, 1010
long-term effectiveness against eye gnat, 785
persistence of in soils, 141
resistance of mosquito to, 777
solubilization of by lipoprotein, 34
Dieldrin, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
salt-marsh caterpillar, 78
Dieldrin, effect of on
apple, 487
boll weevil, 294
corn flea beetle, 822
Cyclocephala spp., 822
Japanese beetle, 821
lesser cornstalk borer, 665

mite nutrition, 487
Phyllophaga spp., 822
rootworms, 822
Sacadodes, 294
seed-corn maggot, 1016
sweet potato flea beetles, 822
toothed flea beetles, 822
Dieldrin, for control of
army cutworm, 319
bean leaf skeletonizer, 566
boll weevil, 229, 1051
corn root webworm, 670
cotton fleahopper, 1051
cranberry fruitworm, 1118
currant fruit weevil, 475
cutworms, 526
desert grasshopper, 724
European pine shoot moth, 366
eye gnats, 362
fuller rose beetle, 164
Hyperodes humilis, 255
imported fire ant, 190
lygus bugs, 397
mint insects, 527, 529
mint looper, 527
Nantucket pine moth, 366
plum curculio, 439
spider mites, 1053
sugar-beet wireworm, 177
three-lined potato beetle, 480
thrips, 1052
tobacco flea beetle, 963
woolly apple aphid, 217
Dieldrin, residues of in/on
alfalfa-brome grass pasture, 412
forage, 821
meat, 425
milk, 425
mint oil, 527
radish, 832
soil, 194
sugar beet, 833
Dieldrin, toxicity of to
Chrysops, 263
corn earworm, 22
house fly, 888
livestock, 425
Oriental fruit moth, 470
Dieldrin-Risella, susceptibility to of, Anopheles mosquitoes, 102
Diel behavior, of alfalfa weevil, 675
Diet
artificial, growth of cigarette beetle on, 382
casein, for laboratory culture of pink bollworm, 791
cottonseed meal, for laboratory culture of pink bollworm, 791
for rearing stable fly, 409
O,O-diethyl-*O*-benzhydryl thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-diethyl-*O*-(4-chloro-propylbenzyl) thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-diethyl-*O*-cinnamyl thiophosphate, toxicity of to Oriental fruit moth, 470
O,O,-Diethyl-*O*-hydrocinnamyl thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-Diethyl-*O*-(*p*-methoxybenzyl) thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-Diethyl *O*-Naphthalimido Phosphorothioate, biological degradation of, 848
O,O-Diethyl-*O*-(α-phenyl propyl) piperonyl thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-Diethyl-*O*-(α-propyl benzyl)thiophosphate, toxicity of to Oriental fruit moth, 470
O,O-Diethyl-*O*-(α-propyl-p-methoxybenzyl) thiophosphate, toxicity of to Oriental fruit moth, 470
Diethyltin diacetate, against house fly, 447
Diethyltin dichloride, against house fly, 447
Diethyltoluamide
deleted common name of insecticide, 677
toxicity of to northern fowl mite, 12
O,O-Diethyl-*O*-(o-veratryl)thiophosphate, toxicity of to Oriental fruit moth, 470
Dilan, effectiveness of on salt-marsh caterpillar, 78
Dilan, for control of
Mexican bean beetle, 819
mint insect, 529
salt-marsh caterpillar, 76
Dilan, toxicity of to
corn earworm, 22
Oriental fruit moth, 470
vedalia beetle, 1056
Dimetan, stability of, 205
Dimethoate
against green peach aphid, 624
against European corn borer, 625
effect of on survival of surpentine-leaf-miner parasite, 872
honey bee mortality from, 1014
residual effectiveness of, 478
resistance of mosquito to, 777
Dimethoate, effectiveness of against
bed bug, 6
Gulf Coast tick, 6
Dimethoate, effect of on
beetle, 1035
plant emergence, 226
Dimethoate, for control of
azalea lace bug, 408, 839
bean leaf skeletonizer, 568
birch leaf miner, 407
boll weevil, 1052
boxwood leaf miner, 407
cattle grubs, 183, 541, 683
clover aphid, 1014
corn earworm, 819
cotton fleahopper, 1052
cranberry fruitworm, 1118
eastern spruce gall aphid, 979
European pine shoot moth, 349
face fly, 450
grape leaf folder, 531, 533
green peach aphid, 871, 963, 1099
horn fly, 477
house fly, 315
Hyperodes humilis, 255
Mexican bean beetle, 819
mites on apple, 736
Oriental rat flea, 167
pea aphid, 819
potato tuberworm, 868
poultry lice and mites, 161
San Jose scale, 1085
sap beetles, 175
screw-worm, 1059
serpentine-leaf-miner, 872
tarnished plant bug, 690
thrips, 227, 1051
two-spotted spider mite, 226
Dimethoate, toxicity of to
cockroach, 25
horses, 602
house fly, 6, 25, 741
mice, 25
white rat, 6
2,5-dimethoxy-*N*-methylcarbamate, against
citrus red mite, 828
house fly, 828
mosquito, 828
salt-marsh caterpillar, 828
3,5-dimethoxyphenyl-*N*-methylcarbamate, against
citrus red mite, 828
house fly, 828
mosquito, 828
salt-marsh caterpillar, 828
m-dimethylamino-*N*-methylcarbamate, toxicity of to salt-marsh caterpillar, 1022
4,4′ dimethylbenzhydryl amine HCl, for control of cattle grubs, 183
Dimethyl benzyl, toxicity of to house fly, 742
3,4-Dimethylbenzyl chrysanthemumat, toxicity of to chicken body louse, 12
Dimethyl divinyltin, against house fly, 447
Dimethyltin dichloride, against house fly, 447
Dimethyl vinyltin-α-bromopropionate, against house fly, 447
Dimethyl vinyltin chloride, against house fly, 447
Dimetilan, for control of
house fly, 315, 898
Dimetilan, toxicity of to
German cockroach, 1008
house fly, 233, 741, 899, 1005
milkweed bug, 1007
rats, 233
Dimite, for control of myocoptic mange mite, 168
Di-*n*-octyltin dichloride, against house fly, 447
Di-*n*-octyltin dilaurate, against house fly, 447
Diphenylacetonitrile, effect of on house fly development, 803
Diphenyltin dichloride, against house fly, 447
Diprion similis, 648
Dipterex
duration of toxicity of to fuller rose beetle, 165
resistance of mosquito to, 777
Dipterex, for control of
cattle grubs, 683
Drosophila, 1109
green peach aphid, 1099
house fly, 899
imported fire ant, 190
lone star tick, 953
northern fowl mite, 4
plum curculio, 335
Dipterex, toxicity of to
Chrysops, 263
corn earworm, 22
house fly, 429, 741, 1066
oriental fruit moth, 469
Disonycha triangularis, 132
Di-Syston
as soil treatment against eye gnat, 1102
honey bee mortality from, 1014
Di-Syston, effect of on
beetle, 1035
germination of wheat, 575
Ladino clover, 1134
lesser cornstalk borer, 665
plant emergence, 226
plants, 838
Di-Syston, for control of
azalea lace bugs, 839
clover aphid, 1014
greenbug, 278
green peach aphid, 725, 835
Hessian fly, 501
hollyscale, 406
leafhopper, 835

lygus bugs, 402, 767
southern garden leafhopper, 725
spider mites, 725
thrips, 225
tobacco flea beetle, 835
two-spotted spider mite, 224
Dition, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Divinyl-*n*-butyltin chloride, against house fly, 447
Divinyl phenyltin fluoride, against house fly, 447
DMBC, toxicity of to house fly, 741
Dogs, effect of dieldrin on, 826
Dosimetry
for radiological studies of *Tribolium castaneum*, 496
values for biological parameters, 499
Dow 9-B, effect of on germination of wheat, 575
Dow ET-15
resistance of mosquito to, 777
systemic and topical activity of, 692
toxicity of to house fly, 429
toxicity of to Oriental fruit moth, 469
Dow M-350, toxicity of to, corn earworm, 22
Dowco 105, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Dowco 105, for control of cattle grubs, 683
Dowco 105, toxicity of to horses, 602
Dowco 109, for control of
cattle grubs, 183, 541, 683
Oriental rat flea, 167
poultry lice and mites, 161
Dowco 109, toxicity of to horses, 602
Draeculacephala spp., 823
Drosophila, 213, 969, 970
busckii, 8
hydei, 8
melanogaster, 8, 193, 337, 1107
simulans, 969
transversa, 8
Dust-Bath-Boxes, malathion-treated, for control of northern fowl mite, 701
Dylox
as soil treatment against eye gnat, 1102
effectiveness of on salt-marsh caterpillar, 78
residues of on alfalfa hay, 77
Dylox, effect of on
cabbage looper, 77
omnivorous leafroller, 77
southern garden leafhopper, 76
survival of serpentine-leaf-miner parasite, 872
Dylox, for control of
alfalfa caterpillar, 73
corn earworm, 819
cotton leaf perforator, 76
cranberry fruitworm, 1118
Eushistus sp, 75
insects on peas, snap and lima beans, 819
lygus bugs, 74, 767
morning-glory leaf miner, 1119
salt-marsh caterpillar, 76
sap beetles, 175
Say stink bug, 75
tarnished plant bug, 690
variegated cutworm, 74
Dylox, toxicity of to
coccinellids, 68
Nabis ferus, 68
Praon palitans, 68
Orius sp., 68
Chrysopa, spp., 68
Geocoris spp., 68
Dysmicoccus
brevipes, 296, 322
neobrevipes, 322

Economic importance, of chicken body louse, 421
Ectopria, 609
Elasmopalpus lignosellus, 664
Elm, systemic insecticidal action in cortical tissues of, 1034
Empoasca, 823
fabae, 113, 389, 527, 818
filamenta, 835
solana, 76, 725
Emulphor EL 719, in phytotoxic formulations, 203
EN 18133, as soil treatment against eye gnat, 1105
Endrin
duration of toxicity of to fuller rose beetle, 165
effectiveness of on salt-marsh caterpillar, 78
honey bee mortality from, 1014
long-term effectiveness of against eye gnat, 785
loss of from soil surface, 650
susceptibility of boll weevil to, 311
Endrin, against
boll weevil, 711
eye gnat, 367
European corn borer, 624
green peach aphid, 624
Endrin, effect of on
boll weevil, 294
hornworms, 976
Sacadodes, 294
transamination in cockroach, 167
Endrin, for control of
army cutworm, 319
boll weevil, 229
clover aphid, 1014
corn earworm, 397
Douglas-fir beetle, 949
European pine shoot moth, 366
fuller rose beetle, 164
grape leaf folder, 531, 532
green peach aphid, 727, 871, 1099
Hyperodes humilis, 255
imported fire ant, 190
lesser cornstalk borer, 664
lygus bugs, 397
Nantucket pine moth, 366
plum curculio, 439
potato tuberworm, 868
sugar-beet wireworm, 177
sugarcane borer, 935
three-lined potato beetle, 480
Endrin, residual effect of on
bollworm, 104
cabbage looper, 104
cotton leafworm, 104
salt-marsh caterpillar, 104
Endrin, toxicity of to
cabbage looper, 991
Chrysops, 263
corn earworm, 22
Oriental fruit moth, 470
vedalia beetle, 1056
Enoplognatha sp., 812
Entomobryidae, 823
Ephedrus nitidus, 659
Ephemerella, 609
Ephestia
figulilella, 970, 975
kuhniella, 537, 971
Epilachna varivestis, 132, 389, 818
Epitrix, 822
cucumeris, 527
fuscula, 527
hirtipennis, 132, 835, 963
EPN
resistance of mosquito to, 777
toxicity of to Oriental fruit moth, 469
EPN, effect of on
seed-corn maggot, 1016
survival of serpentine-leaf-miner parasite, 872
EPN, for control of
cranberry fruitworm, 1118
serpentine-leaf-miner, 874
EPN, residues of on
forage under pecan, 719
grapes, 441
EPN-300
for control of mint insects, 528
residues of in mint oil, 531
1,2-Epoxyhexadecane, as attractant to gypsy moth males, 173
Equipment
an effective, economical flytrap, 1131
automatic microapplicator, 853
dimetilan baits, fly ribbons and cords, 898
improved trap for deciduous tree fruit flies, 967
trap to recover dead and abnormal honey bees, 782
ventilation unit for aerating insectary sleeve cages, 959
wide-range variable temperature cabinet, 1030
Eridantes erigonoides, 812
Erigonidae, in apple orchards, 811
Erinnys ello, 1128
Eriosoma lanigerum, 217
Erythroneura elegantula, 531, 532
Estigmene, 294
acrea, 76, 78, 104, 294, 828, 957, 1021
Ethion
against desert spider mite, 748
as soil treatment against eye gnat, 1102
for bee poisoning, 1013
honey bee mortality from, 1014
Ethion, for control of
clover aphid, 1014
corn earworm, 402, 819
cotton aphid, 747
cranberry fruitworm, 1118
European pine shoot moth, 352
grape mealybug, 357
lygus bugs, 402, 767
Mexican bean beetle, 819
San Jose scale, 1085
screw-worm, 1059
serpentine-leaf-miner, 872
southern garden leafhopper, 725
spider mite complex, 522
spider mites, 725
tarnished plant bug, 690
Ethion, toxicity of to
corn earworm, 22
Oriental fruit moth, 469
serpentine-leaf-miner parasite, 872
m-ethoxy-*N*-methylcarbamate
against citrus red mite, 828
against house fly, 828
against mosquito, 828
against salt-marsh caterpillar, 828
toxicity of to salt-marsh caterpillar, 1022
O-ethoxy-*N*-methylcarbamate
against citrus red mite, 828
against house fly, 828
against mosquito, 828
against salt-marsh caterpillar, 828

toxicity of to salt-marsh caterpillar, 1022
Ethylene chlorobromide, susceptibility of Mexican Fruit Fly to fumigation with, 323
Ethylene dibromide
distribution and sorption of applied to wheat by recirculation, 259
effect of on confused floor beetle, 762
for control of stored-grain insects, 536
fumigation of buildings with, 508
Ethylene dibromide, residues of on
blackeye bean, 833
lettuce, 833
lima bean, 832
onion, 832
spinach, 833
sugar beet, 832
table beet, 832
turnip, 832
Ethylene dichloride,
distribution and sorption of applied to wheat by recirculation, 259
for control of stored-grain insects, 536
O-Ethyl *S*-2(ethylsulfinyl)-ethyl *n*-propylphosphonothiolate, as systemic insecticide, 127
Ethyl Guthion
as soil treatment against eye gnat, 1102
for control of codling moth, 1083
N(2-Ethylhexyl)bicyclo-2,2,1-hept-5-ene-2,3-dicarboxamide, screening of as cockroach repellent, 806
Eucosma gloriola, 248
Eugenol, effect of on house fly development, 803
Eulasonia sp., 1097
Eupelmus, 305
cyaniceps cyaniceps, 305
limneriae, 305
Eurytoma, 537
appendigaster, 305
Euschistus sp., 75
variolarius, 527
Eustala anastera, 813
Euzophera semifuneralis, 276
Experiments, designing of for European mite control in orchards, 892
Experimental Nematocide 18133, as soil treatment against eye gnat, 1102

Fannia canicularis, 999
Fat baits, for control of imported fire ant, 188
Feltia, 294
ducens, 319
Fenusa pusilla, 406
Fertilizer, effect of on systemic insecticide effectiveness against greenbug, 278
Fir, balsam, as overwintering site of spruce budworm, 684
Fish, bluegill, toxicity of imported fire ant to, 1137
Flax, effect of fumigants on germination of, 17
Flight distance, of boll weevil, 681
Flight range, of European chafer, 222
5-Fluorouracil, effect of on house fly development and fertility, 803
Fly ribbons, for house fly control, 898
Flytrap, an effective, economical, 1131
Food, adult, effect of on susceptibility of boll weevil, 534
Foraging behavior, of honey bee before and after flowering, 564
Forbs, grasshopper damage to, 640
Frankliniella, 1051
fusca, 131
gilmorei, 527
moultoni, 224
Frontinella communis, 812
Fruit, dried, experiments to exclude *Drosophila* from, 969
Fruit drop, caused by Sevin, 956
Fumarin, in systemic insecticide baits for rat flea control, 169
Fumigants, liquid, distribution and sorption of applied to
wheat by recirculation, 259
Fumigant Vaporizer Unit, for laboratory use, 562
Fumigation
for control of dry-wood termite, 503
reponse of quiescent Khapra Beetle larvae to, 698
Fungi, storage, development of in wheat, 375
Fungus, parasitic on seed-corn maggot, 478
Fungus disease, of salt-marsh caterpillar, 957

G-30494
as soil treatment against eye gnat, 1102
toxicity to house fly, 741
GC-1189, against eye gnat, 367
GS-3582, as soil treatment against eye gnat, 1102
GC-3583, as soil treatment against eye gnat, 1102
GC-4072, as soil treatment against eye gnat, 1102
Gambrus ultimus, 305
Gamtox, for control of
American plum borer, 276
pear borer, 276
Gasterophilidae, 602
Geigy G-30494
against boll weevil, 747
against boll worm, 747
against desert spider mite, 748
for control of cotton aphid, 747
Gelis, 305
tenellus, 305
General Chemical 3583, for control of screw-worm, 1059
General Chemical 4072, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
General Chemical 4072, for control of screw-worm, 1059
Genetics, of parathion resistance in Pacific spider mite, 626
Geocoris, 68
punctipes, 1088
Germination, effect on of
phorate in sorghum, 154
virus yellows and curly top in sugar beet, 164
Glischrochilus spp., 823
Glyodin, for control of European red mite, 892
Glypta sp., 251
Gnorimoschema operculella, 868
Goat, malathion residues in tissues of, 972
Goldenrod, as forage for bees, 679
Goniocotes gallinae, 328
gigas, 328
Grain, stored, relation of moisture and temperature of to
effectiveness of fumigants on under forced circulation, 1071
Graminella nigrifrons, 823
Grammonota inornata, 810
Grape, residues of Sevin on, 858
Grapholitha molesta, 304, 468
Grasses,
desert grasshoppers on, 724
grasshopper damage to, 640
residues of DDT on, 1020
residues of 1-Naphthol on, 484
residues of Sevin on, 484
Grasshoppers, long-horned, control of on mint, 527
Greens, mustard, residues of nicotine on, 64
Gryllidae, 823
Gryllotalpa africana, 1018
Guinea pigs, for insecticide screening tests, 1125
Guthion
as soil treatment against eye gnat, 1102
duration of toxicity of to fuller rose beetle, 165
for bee poisoning, 1013
residual value against plum curculio, 335
resuidues of on forage under pecan, 719
seasonal tolerance of boll weevil to, 41
susceptibility of boll weevil to, 311
Guthion, against
boll weevil, 711, 747
bollworm, 711, 747
European corn borer, 624
green peach aphid, 624
Guthion, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 52
desert spider mite, 54
Guthion, effect of on
boll weevil, 294
lesser cornstalk borer, 665
plant emergence, 226
Sacadodes, 294
survival of serpentine-leaf-miner parasite, 873
Guthion, for control of
bean leaf skeletonizer, 568
boll weevil, 228, 1051
clover aphid, 1013
codling moth, 1083
cotton fleahopper, 1051
cotton leaf perforator, 76
cranberry fruitworm, 1118
Drosophila, 1109
European pine shoot moth, 349, 365
fuller rose beetle, 164
grape berry moth, 857
green peach aphid, 871, 1099
Hyperodes humilis, 251
imported fire ant, 190
lygus bugs, 402, 514
Nantucket pine moth, 365
Oligonychus pratensis, 1061
potato tuberworm, 868
San Jose scale, 1085
serpentine-leaf-miner, 873
spider mites, 1053
spider mite complex, 522
tarnished plant bug, 690
thrips, 225
two-spotted spider mite, 225, 967
woolly apple aphid, 217
Guthion, toxicity of to
boll weevil, 534
Chrysops, 263
corn earworm, 22
Oriental fruit moth, 469

Habrobracon juglandis, 971
Habrocytus cerealellae, 537
Habrophlebia, 609
Hadrotettix trifasciatus, 641

Haematopinus eurysternus, 904
Haemolaelops glasgowi, 904
Haltichella xanticles, 305
Halticoptera aenea, 682
Heliothis zea, 22, 54, 76, 104, 118, 127, 131, 294, 318, 389, 397, 527, 695, 711, 745, 747, 818, 823, 849, 1051, 1088
Henoticus sp., 537
Hentzia palmarum, 812
Heptachlor
against eye gnat, 367
epoxidation of in soils, 192
for protection of crated furniture, 966
solubilization of by lipoprotein, 34
Heptachlor, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
Heptachlor, effect of on
flavor of carrots, parsnips and turnips, 285
Ladino clover, 1134
lesser cornstalk border, 665
predators and parasites of root maggots, 932
seed-corn maggot, 1016
survival of serpentine-leaf-miner parasite, 874
turnip maggot, 1017
Heptachlor, for control of
alfalfa caterpillar, 73
black carpet beetle, 966
boll weevil, 229
clover root borer, 449
corn root webworm, 670
cutworms, 526
desert grasshopper, 724
Douglas-fir beetle, 949
European pine shoot moth, 366
eye gnats, 362
Hyperodes humilis, 253
imported fire ant, 188, 647
Japanese beetle, 320
lygus bugs, 398
mint insects, 527
mint looper, 527
Nantucket pine moth, 366
plum curculio, 335, 439
root maggots, 65, 932
sand wireworm, 964
serpentine-leaf-miner, 874
sugar-beet wireworm, 177
tobacco flea beetle, 963
variegated cutworm, 74
Heptachlor, residues of in/on
alfalfa-brome grass pasture, 412
corn stover, 19
mint oil, 527
soil, 194
Heptachlor, toxicity of to
Chrysopa spp., 70
Chrysops, 263
Collops vittatus, 70
convergent lady beetle, 70
corn earworm, 22
Geocoris spp., 70
house fly, 741
Nabis ferus, 70
Oriental fruit moth, 470
Orius sp., 70
Heptachlor epoxide
analysis for in milk, 19
excretion of in milk of cows fed on heptachor-treated pastures, 411
formed from heptachlor in soil, 192
residues of in soil, 194
toxicity of to house fly, 741
Hercothrips phaseoli, 105
Hercules 1986, toxicity of to Oriental fruit moth, 470
Hercules 2032, toxicity of to Oriental fruit moth, 470
Hercules 2118, toxicity of to Oriental fruit moth, 470
Hercules 3273, toxicity of to Oriental fruit moth, 470
Hercules AC-3895, toxicity of to Oriental fruit moth, 469
Hercules 3895, systemic effectiveness of to bed bug, 6
Gulf Coast tick, 6
Hercules 3895, toxicity of to
house fly, 6
white rat, 6
Hercules 5727
against boll weevil, 712
against bollworm, 712
for control of sand wireworm, 964
Heterococcus graminicola, 737
Hexa-*n*-butyl distannane, against house fly, 447
Hexachlorocyclopentadiene, for control of grape phylloxera, 788
5-Hexenoic acid, as synergist in house fly sprays, 300
Hippelates
collusor, 362, 367, 650, 785, 1102
hermsi, 362
Hippodamia, 767
convergens, 69, 91, 257
g. glacialis, 91
parenthesis, 91
quinquesignata punctulata, 68
5-signata ambigua, 659
Histopathological effects, of cattle grub on gullet tissue, 280
Hog
malathion residues in tissues of, 972
residues of toxaphene in, 322
Honey bee, abundance of, 679
Horistonotus uhlerii, 963
Horogenes sp., 1097
Horse, toxicity to of
DDVP, 593
organic phosphorus insecticides, 602
Horse meat, for mass-production of screwworms, 1114
Humidity, relative, effect of on diapuase in pink bollworms, 796
Hydrocyanic acid, effect of on
confused flour beetle, 1071
flax seed germination, 17
rice weevil, 1071
Hydrocyanic acid, fuminagion of khapra beetle larvae with, 698
Hydrodarco B, effect of on
plant sugars, 491
two-spotted spider mite, 492
Hydrogen phosphide, effect of on seed germination, 1
Hydroptila, 609
2-Hydroxyethyl n-octyl sulfide, screening of as cockroach repellent, 806
Hylastinus obscurus, 865, 449
Hylemya
antiqua, 1015
brassicae, 65, 932
ciliicrura, 65, 227, 478, 934, 1015
floralis, 1015
fugax, 55
liturata, 65
platura, 1015
Hypera
nigrirostris, 823
postica, 178, 570, 675
punctata, 823
Hyperaphis
binotata, 325
signata, 325
Hyperodes humilis, 251
Hypoderma, 125, 541, 1125
bovis, 183, 339, 683, 729, 814, 1126
lineatum, 183, 280, 339, 683, 729, 814, 1125

Icerya purchasi, 1055
Idechthis canescens, 537
Indalone, repellency of to stable fly, 434
Indole butyric acid, effect of on house fly development, 803
Infestation, of imported fire ant, 646
Infestation criteria, for determining severity of by clover aphid, 1015
Insectary, an inexpensive indoor type, 685
Insectary production, of *Stethorus* spp., 982
Insect generations, change in importance of segments of, 694
Insecticidal baits, for control of imported fire ant, 188
Insecticides
analysis of joint action of against house fly, 887
apparatus for assay of, 483
differential toxicity of to cockroach, 25
differential toxicity of to mice, 25
differential toxicity of to house fly, 25
differential toxicity of to insects, 25
that change the importance of segments of insect generations, 694
Insecticides, carbamate
as soil treatments against eye gnat, 1102
methylenedioxyphenyl derivatives as synergists for, 231
reaction mechanisms of with esterases, 205
stability of, 205
systemic properties of, 1021
toxicity of, 205
Insecticides, chlorinated hydrocrabon
as soil treatments against eye gnat, 367, 1102
loss of activity of in soil, 785
loss of from soil surface, 650
resistance of mosquitoes to, 1122
stability of resistance in mosquitoes, 1122
Insecticides, granulated
efficiency of with solvents used for impregnation, 938
Insecticides, organophosphorus
as soil treatments against eye gnat, 1102
house fly resistance to, 313
resistance of mosquito to, 777, 1122
stability of resistance to in mosquitoes, 1123
toxicity of to horses, 602
Insecticides, phosphorothioate systemic and topical activity of, 692
Insecticides, silica gel susceptibility of mosquitoes to, 915
Insecticides, soil
effect of on apple trees, 487
effect of on mite nutrition, 487, 491
effect of on plant sugar, 491
Insecticides, substituted phosphate
systemic and topical activity of, 692
Insecticides, systemic
against cattle grubs, 183
alkyl-phosphonate esters as, 127
as feed additives for cattle grub control, 682
effect of on germination of wheat, 575
in vivo activity of, 183

preliminary evaluation of several, 1125
effect of acceleration force on mortality of, 691
Insects,
apple, control of, 685
aquatic, effect of DDT on, 608
on trees that develop oakwilt, 578
soil, control of in Japan, 1015
surveys of in central Rocky Mountain forests, 458
use of for own destruction, 415
Isia isabella, 527
Isobutyric acid, as synergist for barthrin in sprays
against house fly, 284
Isodrin, for control of
lygus bugs, 398
plum curculio, 439
Isodrin, toxicity of to
corn earworm, 22
Oriental fruit moth, 470
Isogenus, 609
Isolan, stability of, 205
Isolan, toxicity of to
German cockraoch, 1008
house fly, 206, 233, 1005
milkweed bug, 1007
mouse, 206
rats, 233
salt-marsh caterpillar, 1022
Isoperla, 609
o-isopropoxy-*N*-methylcarbamate, against
citrus red mite, 828
house fly, 828
mosquito, 828
salt-marsh caterpillar, 828
m-isopropyl-*N*-methylcarbamate, toxicity of to salt-marsh caterpillar, 1022
Isotox, for control of
American plum borer, 276
pear borer, 276
Itoplectis conquisitor, 305

Japygidae, 823

Kale, residues of nicotine, on, 64
Karathane, influence of an apple growth, 717
Kelthane, duration of toxicity of to fuller rose beetle, 165
Kelthane, effect of on
demeton-resistant citrus red mites, 388
ovex-resistant citrus red mites, 388
Kelthane, for control of
European red mite, 685
grape leaf folder, 531, 533
cotton mite, 204
mites, 735
Oligonychus pratensis, 1061
spider mites, 725
spider mite complex, 522
Kepone
against European corn borer, 625
toxicity of to corn earworm, 22
Kepone, for control of
bean leaf skeletonizer, 566
codling moth, 1083
European pine shoot moth, 366
grape leaf folder, 531, 533
green peach aphid, 404, 1099
imported fire ant, 188, 190
Nantucket pine moth, 366
Korlan (*see Ronnel*)

Lachesilla pedicularia, 537
Lacon binodulus, 1018
Laemophloeus
ferrugineus, 536
pusillus, 537
Laetilia coccidivora, 325
Lanthus, 609
Laphygma
exigua, 73
frugiperda, 319, 389
Lariophagus distinguendus, 537
Lasioderma serricone, 381
Lathridius minutus, 537
Lead arsenate, effect of on parasitization of Oriental fruit moth cocoons, 305
Lead arsenate, for control of
grape leaf folder, 532
mites on apple, 735
Leafminers, control of on mint, 527
Leaf tier, control of on mint, 527
Lema trilineata, 480, 527
Lepidoptera, attracted to black light, 173
Lepidostoma, 609
Lethane, for control of
European pine shoot moth, 366
Nantucket pine moth, 366
Lethane 384, toxicity of to *Chrysops*, 263
l-(—)-Leucine, effect of on house fly development, 803
Lime, effect of soil application on Japanese beetle, 476
Lime sulfur, for control of San Jose scale, 1086
Limonius californicus, 176
Lindane
effect of on quality and yield of mushrooms, 558
for protection of crated furniture, 966
influence on growth of apple, 717
long-term effectiveness against eye gnat, 785
persistence of in soils, 136
persistence of residues on alfalfa, 960
residual studies of, 193, 196
resistance of house fly to, 313
solubilization of by lipoprotein, 34
Lindane, against
eye gnat, 367
mosquitoes, 124
raisin moth, 975
Lindane, for control of
apple aphid, 718
black carpet beetle, 966
Douglas-fir beetle, 949
eastern spruce gall aphid, 979
imported fire ant, 190
lone star tick 953
mint insects, 529
potato leafhopper, 718
woolly apple aphid, 217
Lindane, toxicity of to
Chrysops, 263
corn earworm, 22
house fly, 889
Linkage groups, of house fly, 985
Lipeurus caponis, 160, 328
Lipoprotein, solubilization of insecticides by, 31
Liriomyza
munca, 872
pictella, 682
Lissorhoptrus oryzophilus, 1135
Litargus dexpunctatus, 537
Locustidae, on Ladino clover, 114
Longitarsus
suspectus, 527
waterhousei, 526, 527
Loxostege, 73
similalis, 527
sticticalis, 585
Lucerne, honey bee count on, 565
Lyctocoris campestris, 537
Lydella grisescens, 582, 824
Lygus, 75
elisus, 689
hesperus, 327, 397, 689, 767
lineolaris, 132, 513, 527, 689, 690, 823, 1063
Lymeon orbum, 305

Macrodactylus subspinosus, 527
Macrosiphum
gaurae, 258
granarium, 197
pisa, 113, 151, 235, 258, 655, 818, 823, 863, 881
rosae, 151, 258
solanifoli, 389, 655, 868
Macrosteles, 823
fascifrons, 527
Magicicada septendecim, 961
Malacosoma fragile, 461
Malaoxon, resistance of mosquito to, 777
Malathion
as grain protectant, 341
development of resistance to in cockroach, 1143
duration of toxicity to fuller rose beetle, 165
effect of on soil microarthropods, 357
effect of on transamination in cockroach, 167
effect of on soybean yield, 133
for reducing dried-fruit packing house infestations, 969
in antagonism-penetration studies of house fly, 14
influence of on apple growth, 717
interference of in biological control of cottony-cushion scale, 1055
residual effectiveness of, 478
resistance of mosquito to, 777, 1122
susceptibility of green peach aphid to, 86
tolerance of green peach aphid to, 87
Malathion, against
boll weevil, 711
lead cable borers, 968
mosquitoes, 124
raisin moth, 975
stable fly, 303
Malathion, effectiveness of on
boll weevil, 53
bollworm, 54
cotton fleahopper, 53
desert spider mite, 54
Malathion, for control of
apple aphid, 685, 718
apple seed chalcid, 166
bean leaf skeletonizer, 567
beet armyworm, 618
boll weevil, 228
chicken lice, 328
chicken mite, 328
cockroach, 200
codling moth, 1084
cotton leaf perforator, 76
cranberry fruitworm, 1116
Dropsophila, 1107
eastern spruce gall aphid, 979
European corn borer, 590
European pine shoot moth, 365
face fly, 450
grape leaf folder, 532
grape mealybug, 356
horn fly, 477
house fly, 315
fire ant, 190
leafhoppers, 527
lone star tick, 953
lygus bugs, 74, 514
mint plant bugs, 527
morning-glory leaf miner, 1119

Nantucket pine moth, 365
northern fowl mite, 701
Oligonychus pratensis, 1061
pickleworm, 482
plum curculio, 335
potato leafhopper, 718
poultry lice and mites, 160
San Jose scale, 1085
sap beetles, 175
screw-worm, 1059
spider mite, 527
spider mite complex, 522
stored-grain insects, 536
Malathion, residues of on/in
alfalfa, 495
bean leaves, 452
cucumbers, 483
meat, 495
milk, 495
tissues of sheep, goats and hogs, 972
Malathion, toxicity of to
chicken body louse, 12
chicken mite, 12
Chrysops, 263
northern fowl mite, 12
corn earworm, 22
house fly, 429, 899
Oriental fruit moth, 470
vedalia beetle, 1056
Malathion, P^{32}-labeled, rate of detoxification of mosquito, 779
Malathion, diethyl homolog of, resistance of mosquito to, 779
Malathion resistance, effect of DDT resistance on development of, 320
Malathion, SF59, residual effectiveness of, 478
Mangora placida, 813
Marking, mass, of foraging honey bees, 696
Mass production
methods for colonization of *Culicoides*, 731
of citrus red mite, 964
of screw-worms, 1110
Mastrua pilifrons, 305
Meat, residues of malathion in, 495
Medium
for aseptic rearing of stable fly, 408
for rearing some lepidopterus corn insects, 318
fly, effect on house fly of adding wheat germ and corn oil to, 325
Megaselia agarici, 558
Melanophila drummondi, 951
Melanoplus, 823
bilituratus, 640
cuneatus, 640
differentialis, 132, 527
femur-rubrum, 527
Melonotus, 527
caudex, 1018
legatus, 1018
Membracidae, attracted to black light, 173
Membrane, animal, use of in evaluation of repellents against stable fly, 432
Menacanthus stramineus, 12, 160, 328, 421
Menopon gallinae, 160, 328
Mermithidae, as parasites of the alfalfa weevil, 179
Mesochorus perniciosis, 585
Metabolism
effects of on house fly by certain compounds, 802
effect of P^{32}-Delnav in of steer, 60
effect of P^{32}-Delnav in of mice, 62
importance of in determining insecticide toxicity of insects and mice, 25
Metabolites, of P^{32}-Delnav in cattle, 672
Metacide
effect on survival of serpentine-leaf-miner parasite, 873
for control of serpentine leaf miner, 873
Metaphidippus galathea, 813
Meteorus
autographae, 567
loxostege, 585
Methallyl chloride, for control of rice weevils, 288
Methionine, effect of on house fly development, 803
Methods
colorimetric, for determination of Co-Ral in animal tissue, 263
detection of gypsy moth with natural female sex lure, 751
development of synthetic medium for aseptic rearing of stable fly, 408
for calculating joint action of insecticides, 888
for comparing powder-post beetle populations in substructures of buildings, 329
for determining damage by sorghum midge in grain sorghum, 749
for determining infestation of sorghum webworm, 749
for evaluation of stable fly toxicants and repellents, 301
for screening compounds as attractants to gypsy moth males, 172
infrared determination of aldrin and dieldrin in aldrin-treated soil, 1129
insectary production of *Stethorus*, 979
laboratory rearing for *Lygus hesperus*, 327
mass production of citrus red mite, 964
mass production for colonization of mosquitoes, 731
new techniques for screening cockroach repellents, 805
new technique for shipment of natural enemies of insects, 970
sampling experimental plots of alfalfa for insecticide residues, 1078
stalk shredding and plowing for controlling pink bollworm, 436
use of animal membrane in evaluation of chemical repellents, 432
Methoxychlor
against stable fly, 303
residual effectiveness of, 478
solubilization of by lipoprotein, 34
Methoxychlor, for control of
boll weevil, 551
cranberry fruitworm, 1118
European pine shoot moth, 349
horn fly, 477
imported fire ant, 190
mint insects, 529
Methoxychlor, toxicity of to
corn earworm, 22
oriental fruit moth, 469
vedalia beetle, 1056
bis [1(3)-Methyl-3-acetoxy-propyl] adipate, effect of on horse fly development, 803
Methyl bromide
effect of on germination of flax seed, 17
escape of through wallboards, 968
fumigation of buildings with, 503
fumigation of khapra beetle larvae with, 698
fumigation effect on confused flour beetle, 1071
fumigation effect on rice weevil, 1071
N-methylcarbamates
as insecticides, 828
systemic insecticidal properties of, 1021
Methyl-demeton, effect of on *Sacadodes*, 295
3,4-methylenedioxy-*N*-methylcarbamate, against
cirtus red mite, 828
house fly, 828
mosquito, 828
salt-marsh caterpillar, 828
Methylenedioxyphenyl derivatives, as synergists for carbamate insecticides on DDT-resistant and parathion-resistant house flies, 231
5-Methylhydrouracil, effect of on house fly development, 803
Methyl parathion, against
boll weevil, 711
bollworm, 711
cotton leafworm, 749
Methyl parathion, antagonistic action of, 887
Methyl parathion, effect of on
boll weevil, 294
Sacadodes, 294
Methyl parathion, for control of
boll weevil, 228, 1051
cotton fleahopper, 1051
cranberry fruitworm, 1116
European pine shoot moth, 365
Nantucket pine moth, 365
salt-marsh caterpillar, 78
Say stink bug, 75
serpentine-leaf-miner, 874
spider mites, 1053
thrips, 1052
Methyl parathion, toxicity of to
Chrysops, 263
house fly, 889
oriental fruit moth, 469
alpha-(2-Methylphenoxy)-*n*-butyric acid, effect of on house fly development, 803
Methyl trithion
as soil treatment against eye gnat, 1102
honey bee mortality from, 1014
Methyl trithion, for control of
clover aphid, 1012
corn earworm, 819
grape leaf folder, 531, 533
pea aphid, 819
tarnished plant bug, 690
6-Methyluracil, effect of on house fly development, 803
MGK 264, repellency of to
horn fly, 435
stable fly, 435
MGK R-11, toxicity of to corn earworm, 22
MGK R-326, repellency of to
horn fly, 435
stable fly, 434
MGK R-326, toxicity of to corn earworm, 22
Microapplicator, automatic, 853
Microarthropods, effect of insecticides on, 357
Microgramme ruficollis, 537
Microscope, X-ray, to study grain insects, 678
Migration, method of by
greenbug, 605
harlequin bug, 605
Milk, residues in of
DDT, 1020
dieldrin, 827
malathion, 495

Sevin, 327
Millipede, control of on mint, 527
Milo, effect of hydrogen phosphide on seed germination of, 1
Mimetus epeiroides, 813
Misumenops asperatus, 812
Mites
effect of soil insecticides on nutrition of, 487
soil, effect of insecticides on, 358
Mobil GPD 1442, phytotoxicity of, 203
Moisture
effect of on germination and growth of wheat, 771
grain, effect on rice weevil in green corn, 456
grain, influence on effectiveness of malathion, 341
soil, influence on beet webworm, 585
Moisture content, relation to temperature in effectiveness of fumigants in stored grain, 1071
Monacrin
effectiveness of against cattle grubs, 184
for control of cattle grubs, 183
Monarthropalpus buxi, 406
Monocalcium arsenate, effect of on boll weevil, 845
Monsanto CP-7769, against
boll weevil, 711
bollworm, 711
Monsanto CP-7769, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
Monsanto CP-8574
effect of on plant emergence, 226
for control of thrips, 227
for control of two-spotted spider mite, 226
Monsanto CP-10502
effect of on plant emergence, 226
for control of thrips, 227
for control of two-spotted spider mite, 226
Monsanto CP-10502, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Monsanto CP-10516, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Monsanto CP-10561, for control of poultry lice and mites, 161
Monsanto CP-12432, for control of poultry lice and mites, 161
Mormidea lugens, 527
Mortality factors, for vetch bruchid, 555
Mouse, toxicity of insecticides to, 25
Murgantia histrionica, 604
Musca
autumnalis, 450
domestica, 6, 14, 25, 35, 99, 127, 171, 205, 213, 231, 247, 282, 299, 313, 320, 325, 429, 445, 479, 539, 593, 691, 696, 709, 741, 774, 802, 828, 849, 855, 858, 887, 898, 904, 909, 956, 985, 999, 1004, 1037, 1066, 1122, 1133
Mushrooms, from beds treated with lindane, 558
Mutants, house fly, 985
Mutations, in house fly, 213
Mycetophagus, 537
pleuripunctatus, 537
Myocoptes musculinus, 168
Myristyl chloroacetate, 99
Myzus
ascalonicus, 177
cerasi, 659
persicae, 86, 87, 113, 162, 389, 403, 590, 624, 655, 725, 769, 835, 868, 963, 1099

Nabidae, as predators of spotted alfalfa aphid, 92
Nabis ferus, 68, 91, 823, 1088
alpha-Naphthalene acetic acid, effect of on house fly development, 803
Naphthoic acid, as synergist for barthrin in sprays against house fly, 284
1-Naphthol, residues of on forage, 484
alpha-Naphthyl acetamide, effect of on house fly development, 803
Naphthyl methyl isothiourea hydrochloride, effect of on house fly development, 803
beta-Naphthoylacetonitrile, effect of on house fly development, 803
beta-Naphthoxyacetamide, effect of on house fly development, 803
alpha-Naphthyl thiourea, effect of on house fly development, 803
Nasutoideus frontalis, 527
Natural Control, of pine tortoise scale, 325
Nemagon, for control of woolly apple aphid, 217
Nemapogon granella, 537
Nematodes, parasitic on the alfalfa weevil, 179
Nemoura, 609
Neophylax, 609
Neoscona domiciliorum, 813
Nesosteles spp., 823
Nesting habits, of the wasp, *Solierella*, 958
Nesting sites, some characteristics of alkali bee, 424
Neurocolpus nubilus, 527
Nezara viridula, 1128
Nicarbazin
effectiveness of against cattle grubs, 184
for control of cattle grubs, 183
Nicotine, residues of on
cauliflower, 64
celery, 64
green beans, 64
kale, 64
mustard greens, 64
spinach, 64
Nicotinic acid, effect of on house fly development, 803
Nigronia, 609
2-(p-nitrobenzamido)-5-nitrothiazole, for control of cattle grubs, 183
Nitrogen
absorption of in apples treated with soil insecticides, 489
effect of DDT on, 492
effect of Hydrodarco B on, 492
alpha-Nitroso-*beta*-naphthol, effect of on house fly development, 803
Nomia melanderi, 424, 1025
Norvellina seminuda, 527
Nosema apis, 95, 1137
Nosema disease
distribution and severity of infection, 97
in honey bee, 95
seasonal trends of, 97
NPD, in mint insect control, 529
Nuchar C-115-A, as absorbent, effect of on plant sugars and mite nutrition, 491
Nysius ericae, 527

Oak, black and scarlet, insect wounds in, 578
Oak wilt, in black and scarlet oaks, 578
Oats, effect of hydrogen phosphide on seed germination of, 1
Obituary
Clyde Carney Hamilton (1890–1959), 1147
Frederick Zeller Hartzell (1879–1958), 180
Howard Herman Jewett (1884–1959), 181
William Swift Regan (1884–1959), 700
Frank Atwood Sirrine (1861–1959), 700
Otto Herman Swezey (1869–1959), 332
Emilio Viale (1921–1959), 486
Willard Wilson Yates (1888–1960), 1147
Octanamide, as synergist for barthrin in sprays against house fly, 284
Octyl caprylate, as attractant to gypsy moth males, 173
Oil
dormant, for control of American plum borer, 276
dormant, for control of pear borer, 276
dormant, for control of San Jose scale, 1085
emulsified, in copper-zineb sprays for control of citrus rust mites, 395
Mobil, GPD 1432, phytotoxicity of to cotton, 204
Mobil, GPD 1442, phytotoxicity of to cotton, 204
Phillips, base No. 2, phytotoxicity of to cotton, 204
Standard, Chevron Paraffin, phytotoxicity of to cotton, 204
Standard, Pacific base 50 Pale, phytotoxicity of to cotton, 204
Union, No. 90 neutral, phytotoxicity of to cotton, 204
Volck Supreme, for control of San Jose scale, 1085
Oligonychus pratensis, 119, 1061
Olla abdominalis, 91
Oncopeltus fasciatus, 1004
Ophiderma
evelyna, 173
flava, 173
grisea, 173
pubescens, 173
salamandra, 173
Ophionyssus natricis, 903
Opius, 682
suturalis, 682
Organophosphates
animal systemic effectiveness of, 6
for control of Hessian fly, 502
selective toxicity of, 6
Organotin compounds, relationships between structure and insecticidal activity in, 445
Orgilus obscurator, 318
Orius, 69, 725
insidiosus, 91, 584, 823, 1088
tristicolor, 767
Ornithonyssus
bacoti, 475, 904
sylviarum, 4, 12, 160, 476, 701
Orotic acid, effect of on house fly development, 803

Ortho D, for control of
American plum borer, 277
pear borer, 277
pine shoot moth, 352
Oryzaephilus surinamensis, 536, 970
Otobius megnini, 904
Oversummering, of greenbug, 454
Overwintering
of melon leaf miner, 682
of southwestern corn borer, 1128
of spotted alfalfa aphid, 220
Ovex
against two-spotted spider mite, 841
for control of European red mite, 892
for control of a spider mite complex, 522
resistance of citrus red mites to and cross-tolerance to kelthane and tedion, 384
Oxyopes salticus, 813

Pachybrachys
atomarius, 527
othonus, 527
Pachyneuron, 661
siphonophorae, 151
Pagiocerus fiorii, 239
Palatability, of lindane-treated mushrooms, 559
Palorus ratzeburgi, 537
Panogen, effect of on germination of wheat, 575
Panonychus
citri, 384, 828, 964
ulmi, 144, 487, 685, 735, 892
Pantomorus godmani, 164
Papaipema nebris, 527
Paraflow, for control of
American plum borer, 277
pear borer, 277
Paraleptophlebia, 609
Paralobesia viteana, 269, 441, 856
Paraoxon, resistance of mosquito to, 777
Paraphidippus marginatus, 813
Paraphlepsius spp., 823
Para-Scalecide, for control of
American plum borer, 277
pear borer, 277
Parasites
of rose-aphid, 151
of spotted alfalfa aphid, establishment of, 1094
Paratetranychus pratensis, 1061
Parathion
against mosquitoes, 124
as soil treatment against eye gnat, 1103
duration of toxicity of to fuller rose beetle, 165
effectiveness of on salt-marsh caterpillar, 78
interference of in biological control of cottony-cushion scale, 1055
Parathion, effect of on
parasites of oriental fruit moth cocoons and *Trichogramma minutum*, 304
predators and parasites of root maggots, 932
two-spotted spider mite, 81
Parathion, for control of
alfalfa caterpillar, 73
American plum borer, 276
beet armyworm, 618
citrus rust mites, 395
cotton leaf perforator, 76
cranberry fruitworm, 1116
desert grasshopper, 724
European red mite, 892
grape berry moth, 857
grape leaf folder, 531
grape mealybug, 354
green peach aphid, 403, 727
house fly, 898
Hyperodes humilis, 253
imported fire ant, 190
lesser cornstalk borer, 664
lesser peach tree borer, 321
little house fly, 999
lygus bugs, 514
morning-glory leaf miner, 1119
Oligonychus pratensis, 1061
pear borer, 276
plum curculio, 335
root maggots, 932
San Jose scale, 1085
sap beetles, 175
serpentine leaf-miner, 872
spider mite complex, 522
variegated cutworm, 74
Parathion, residues of on
forage under pecan, 719
grapes, 441
Parathion, resistance to of
codling moth, 1083
mosquito, 777, 1122
Pacific spider mite, 626
Parathion, toxicity of to
Chrysopa spp., 68
Chrysops, 263
Coccinellids, 68
cockroach, 25
Collops vittatus, 70
convergent lady beetle, 70
corn earworm, 22
Geocoris spp., 68
Hippodamia parenthesis, 68
Hippodamia quinquesignata punctulata, 68
house fly, 25
mice, 25
Nabis ferus, 68
oriental fruit moth, 469
Orius sp., 68
Praon palitans, 68
serpentine leaf-miner parasite, 872
vedalia beetle, 1056
Peach, honey bee count on, 565
Pear, effect of Sevin on, 956
Peas, uptake of phorate by, 50
Peat moss, as absorbent, effect of on plant sugars and mite nutrition, 491
Pecan, insecticide residues on forage under, 719
Pectinophora gossypiella, 105, 242, 294, 436, 759, 791, 793
Peltoperla, 609
Pentachlorophenyl propionate, toxicity of to oriental fruit moth, 470
6-Pentyl-*m*-tolyl chloroacetate, effect of on house fly development, 803
Peppers, DDT residues on, 622
Perezia pyraustae, 918
Peridroma margaritosa, 74, 527, 823, 849
Periplaneta americana, 25, 166, 170, 171, 808, 904
Perthane
effect of on lesser cornstalk borer, 665
for control of bean leaf skeletonizer, 566
for control of plum curculio, 335
toxicity of to corn earworm, 22
toxicity of to oriental fruit moth, 469
Petrobia latens, 115, 704
Petrova albicapitana, 587
Phaeogenes walshiae walshiae, 305
Phaltan, compatability of with Sevin, 856
Phenacaspis pinifoliae, 406
Phenacoccus solenopsis, 104
Phenococcus solani, 322
Phenothiazine, toxicity of to house fly, 429
1-phenyl-3-(5-nitro-2-thiazolyl) urea, effectiveness of against cattle grub, 184
for control of cattle grubs, 183
Phenyltin trichloride, against house fly, 447
Phidippus
audax, 813
hirsutus, 813
Philaenus leucophthalmus, 527, 823, 960
Philodromus
rufus, 811
vulgaris, 811
Phorate (Thimet)
as soil treatment against eye gnat, 1102
duration of toxicity of to fuller rose beetle, 165
effectiveness of soil injection of against European red mite, 145
effectiveness of soil injection of against two-spotted spider mite, 145
fate of in soils, 47
for bee poisoning in red clover, 1013
germination and growth of wheat seed treated with, 771
honey bee mortality from, 1014
residues of in alfalfa and their effect on cattle, 306
Phorate, effect of on
beetle, 1035
germination of wheat, 575
lesser cornstalk borer, 665
plant emergence, 226
plants, 838
sorghum yield, 154
soybean yield, 133
Phorate, for control of
aphids, 197
apple seed chalcid, 166
azalea lace bugs, 839
birch leaf miner, 407
cabbage aphid, 47
cabbage looper, 47
clover aphid, 1012
corn leaf aphid, 154
green peach aphid, 404, 725, 835, 963, 1099
Hessian fly, 197, 501
holly scale, 406
imported cabbageworm, 47
introduced pine sawfly, 648
leafhopper, 835
lygus bugs, 402
pine needle scale, 406
southern garden leafhopper, 725
spider mites, 725
thrips, 225
tobacco flea beetle, 835
two-spotted spider mite, 224
Phorate, toxicity of to
house fly, 741
oriental fruit moth, 470
Phorodon humuli, 258
Phosdrin
for bee poisoning in red clover, 1013
honey bee mortality from, 1014
residues of on cucumbers, 483
resistance of mosquito to, 777
Phosdrin, for control of
alfalfa caterpillar, 73
alfalfa webworm, 73
bean leaf skeletonizer, 566
beet armyworm, 73
boll weevil, 229
clover aphid, 1013
Drosophila, 1107
eastern spruce gall aphid, 979
European pine shoot moth, 349

green peach aphid, 404
holly scale, 406
Hyperodes humilis, 254
imported fire ant, 190
lygus bugs, 402, 514
Lygus, spp., 74
pickleworm, 482
serpentine leaf miner, 874
two-spotted spider mite, 967
variegated cutworm, 74
Phosdrin, toxicity of to
cabbage looper, 991
Chrysops, 263
house fly, 741
oriental fruit moth, 470
Phosphamidon
against European corn borer, 624
against green peach aphid, 624
as soil treatment against eye gnat, 1102
effect of on beetle, 1035
for bee poisoning in red clover, 1013
honey bee mortality from, 1014
systematic action of in cortical tissues of elm, 1034
Phosphamidon, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Phosphamidon, for control of
azalea lace bug, 408
bean leaf skeletonizer, 568
birch leaf miner, 407
boxwood leaf miner, 407
clover aphid, 1012
corn earworm, 819
cranberry fruitworm, 1118
green peach aphid, 404, 963
Hyperodes humilis, 254
Mexican bean beetle, 819
pea aphid, 819
serpentine leaf miner, 874
P^{32}-Delnav
composition and properties of, 63
effect of on mice, 62
effect of on steer, 60
metabolism of in cattle, 672
residues of on mice, 62
residues of on steer, 60
Phosphorus
absorption of in apples treated with soil insecticides, 489
effect of DDT on, 492
effect of Hydrodarco B on, 492
radioactive, for mass-marking of honey bees, 696
Phostex
honey bee mortality from, 1014
toxicity of to corn earworm, 22
Phostex, for control of
clover aphid, 1014
grape mealybug, 356
Oligonychus pratensis, 1061
San Jose scale, 1086
Photoperiod, effect of on diapause in pink bollworms, 795
Phthia picta, 1128
Phyllocoptruta oleivora, 395
Phyllophaga spp., 527, 822, 860
Phyllotreta
cruciferae, 389
striolata, 389
zimmermanni, 527
Phylloxera vitifoliae, 788
Phytomyza ilicis, 406
Phytophaga destructor, 197, 501, 560, 995
Pieris, 1024
rapae, 47, 389
Pig parlors, control of house fly in, 539
Pimeliaphilus podapolipophagus, 904
Pine
injection of with systemic insecticides, 648
resistance of to pine reproduction weevil, 1044
toxicity of BHC to seedlings of, 175
Pineapple, mealybug populations on in ant-free fields, 296
Piperonal, diethyl acetal as synergist in house fly sprays, 300
Piperonyl butoxide
against raisin moth, 975
as synergist for barthrin in sprays against house fly, 282
as synergist in house fly sprays, 300
effect of on house fly development, 803
penetration of house fly by, 14
synergistic action of, 898
Pirata sp., 813
Pirazinon, toxicity of to oriental fruit moth, 469
Plastanoxus sp., 537
Plathypena scabra, 131, 823
Platynota stultana, 77
Platynychus sp., 1018
Plodia interpunctella, 239, 537, 973
Plowing, for control of pink bollworm, 437
Plutella maculipennis, 389
Plyac, for control of
Hyperodes humilis, 256
San Jose scale, 1086
Podisus
includens, 567
maculiventris, 567
mucronatus, 567
Pollination
alfalfa, effect of apiary distance on, 56
importance of bumble bee vs. honey bee in, 546
red clover, 546
Polyamia inimica, 823
Polybor 3, for control of house fly larvae, 774
Polybutene (Indopol L-100), for control of two-spotted spider mite, 967
Popillia japonica, 320, 476, 582, 691, 821, 860
Population dynamics, in relation to insect control, 416
Porodon menthae, 527
Porthetria dispar, 172, 608, 751, 1019
Potasan
animal systemic activity of, 1037
anticholinesterase activity of, 1039
toxicity of, 1039
Potato
damage to by mole cricket, 1017
damage to by wireworms, 1018
history of aphid control on, 403
uptake of phorate by, 47
Poultry
effect of dieldrin on, 826
young, for control of house fly larvae, 696
Praon
occidentalis, 151
palitans, 68, 151, 235, 655, 1094
unicus, 151
Predator-Prey system, house cricket and house fly, 171
Preimaginal habitat, effect of on susceptibility of boll weevil, 534
Pristomerus appalachianus, 585
Prociphilus fraxinifolii, 258
Prodenia, 294
latifascia, 1128
ornithogalli, 1128
Promoresia, 609
Propane, as synergist for barthrin in sprays against house fly, 284
Propionic acid
as synergist for barthrin in sprays against house fly, 284
as synergist in house fly sprays, 300
m-propoxy-*N*-methylcarbamate, against
citrus red mite, 828
house fly, 828
mosquito, 828
salt-marsh caterpillar, 828
Propyl thiopyrophosphate, toxicity of to oriental fruit moth, 470
Prospaltella perniciosi, 757
Protein baits, for control of imported fire ant, 188
Protoparce
quinquemaculata, 976
sexta, 590, 976, 1128
Psallus seriatus, 52, 1051
Psammotettix spp., 823
Pseudaletia unipuncta, 319, 481
Pseudanthonomus validus, 474
Pseudococcus maritimus, 354, 757
Pseudolimnophilia, 609
Pseudoplusia includens, 566
Pseudoscorpions, effect of insecticides on, 358
Psocids, malathion for reducing dried-fruit packing house infestations of, 970
Psorophora
confinnis, 620
cyanescens, 619
discolor, 620
ferox, 621
horrida, 621
varipes, 621
Psylliodes punctulata, 527
Ptinus
fur, 537
hirtellus, 537
Pupation, of house fly under baseboards, 479
Pyralis farinalis, 537
Pyran, as synergist in house fly sprays, 300
Pyrausta nubilalis, 19, 318, 389, 517, 527, 582, 590, 622, 624, 823, 918
Pyrethrins
as spray against house fly, 299
in aerosols, 909
for control of stored-grain insects, 536
toxicity of to corn earworm, 22
toxicity of to house fly, 889
Pyrethrum, against
raisin moth, 975
stable fly, 303
Pyrolan, stability of, 205
Pyrolan, toxicity of to
German cockroach, 1008
house fly, 206, 233, 1005
milkweed bug, 1007
mouse, 206
rats, 233
Pyrolite, toxicity of to corn earworm, 22

Q-128, toxicity of to corn earworm, 22
Quadraspidiotus perniciosus, 757

R-326, for control of face fly, 450
R-874, as protection against bee poisoning, 1010
Radiation, effect of on *Habrobracon*, 971
Radiobiology, of red flour beetle, 496
Rangeland, short-grass, grasshopper damage on, 640
Raphoplusia ou, 526

Rat
effect of Co-Ral on, 1040
toxicity of DDVP to, 593
Rearing, of spotted alfalfa aphid parasites for mass release, 655
Repellent, as standard term for insect reaction to chemicals, 134
Residues
in established alfalfa treated with granulated phorate, 306
of malathion on and in bean leaves, 452
measuring surface area of apples for expression of, 462
of phorate and their effect on cattle
of Sevin on vegetable crops, 394
Resistance
insecticide, in mosquitoes, 1122
plant, as a means of crop protection, 922
plant, as a means of population control, 923
Rhagoletis
cingulata, 967
cingulata indifferens, 525
fausta, 967
pomonella, 967
Rhipicephalus sanguineus, 904
Rhopalosiphum
fitchii, 113, 197
maidis, 118, 154, 924, 1090
pseudobrassicae, 389
Rhopobota naevana, 1097
Rhyacionia
buoliana, 349, 365
frustrana, 248, 365
Rhyacophila, 609
Rhyzopertha dominica, 537
Riboflavin, effect of on house fly development, 803
Rice, effect of hydrogen phosphide on seed germination of, 1
Roaches, wood, as carriers of oak wilt, 578
Rocket, yellow, as forage for bees, 679
Rodolia
cardinalis, 1055
foedata, 1057
iceryae, 1057
Ronnel (Trolene, Korlan)
against boll weevil, 711
against bollworm, 711
as soil treatment against eye gnat, 1102
effectiveness of against cattle grubs, 184
effect of on rice weevil, 372
effect of on lesser cornstalk borer, 665
residual effectiveness of, 478
resistance to of house fly, 313
resistance to of mosquito, 777
Ronnel, for control of
cattle grubs, 125, 183, 339, 541, 683
Drosophila, 1107
European pine shoot moth, 366
face fly, 450
horn fly, 477
house fly, 315, 539
imported fire ant, 190
lone star tick, 953
Nantucket pine moth, 366
northern fowl mite, 4
oriental rat flea, 167
poultry lice and mites, 161
rat fleas, 169
Ronnell, toxicity of to
chicken body louse, 12
chicken mite, 12
corn earworm, 22
horses, 602
house fly, 429
northern fowl mite, 12
oriental fruit moth, 469
Ronnel, oxygen analog of, resistance of mosquito to, 777
Rotenone
for control of cranberry fruitworm, 1116
for control of insect pests on tomatoes, 1128
solubilization of by lipoprotein, 34
toxicity of to vedalia beetle, 1056
Ruelene
for control of cattle grubs, 814
for control of lone star tick, 953
for control of screw-worm, 1059
toxicity of to horses, 602
Ryania
for control of cranberry fruitworm, 1118
toxicity of to corn earworm, 22
toxicity of to oriental fruit moth, 469

Scadodes pyralis, 291
Salticidae, in apple orchards, 811
Sample location
effect of on rice weevil infestation, 746
effect of on corn earworm infestation, 746
Sampling methods (techniques), for insecticide residues in experimental plots of alfalfa, 1078
for estimating small arthropod populations, 115
for lygus and stink bugs, 689
Sap beetles, as carriers of oak wilt, 578
Schradan, for control of woolly apple aphid, 217
Scobicia declivis, 968
Scolothrips sexmaculatus, 725
Scolytus multistriatus, 1034
Season, influence of on diapause in pink bollworms, 796
Seed abortion, caused by Sevin, 956
Selection pressure, influence of on development of resistance in two-spotted spider mite, 80
Selection response, of parathion resistance in the Pacific spider mite, 626
Senecis jacobaea, 472
Serafume, distribution and sorption of applied to wheat by recirculation, 259
Sericothrips variabilis, 131
Sesamex
against flour beetles, 974
antagonistic action of, 887
as synergist for barthrin in sprays against house fly, 282
synergistic action of, 887
synergistic effect upon mortality of DDT-resistant and parathion-resistant house flies from Sevin, 232
toxicity of to German cockroach, 1008
toxicity of to house fly, 1005
toxicity of to milkweed bug, 1007
Sevin
as soil treatment against eye gnat, 1102
duration of toxicity of to fuller rose beetle, 165
effect of on pear, 956
for bee posoning in red clover, 1013 in protection against bee poisoning, 1010
phytotoxicity of to tomatoes/ 1128
in synergism-penetration studies of house fly, 14
residual effectiveness of, 478
residual value against plum curculio, 335
residue studies of, 326
stability of, 205
Sevin, against
boll weevil, 711, 747
bollworm, 747
cotton leafworm, 747
European corn borer, 624
flour beetles, 974
grape berry moth, 856
Sevin, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
salt-marsh caterpillar, 78
Sevin, for control of
apple seed chalcid, 166
army cutworms, 319
armyworm, 481
bean leaf skeletonizer, 567
beet armyworm, 618
boll weevil, 228, 1051
cabbage flea beetle, 388
cabbage looper, 388
clover aphid, 1013
codling moth, 1083
corn earworm, 388, 402, 819
cotton fleahopper, 1051
cranberry fruitworm, 1116
diamondback moth, 388
Douglas-fir beetle, 949
eastern spruce gall aphid, 979
European corn borer, 388, 590
European pine shoot moth, 349, 366
fall armyworm, 388
grape leaf folder, 531
grape mealybug, 356
green peach aphid, 871, 1099
horn fly, 477
imported cabbageworm, 388
imported fire ant, 190
lone star tick, 953
lygus bugs, 402, 514
Mexican bean beetle, 388
mites on apple, 735
Nantucket pine moth, 366
potato aphid, 388
potato leafhopper, 388
potato tuberworm, 868
poultry lice and mites, 160
San Jose scale, 1085
screw-worm, 1059
southern garden leafhopper, 725
spider mites, 1053
squash bug, 388
tarnished plant bug, 690
thrips, 1051
tomato pests, 1128
turnip aphid, 388
two-spotted mite, 388
woolly apple aphid, 217
Sevin, residues of on
forage, 484
soybeans, 481
vegetable crops, 394
Sevin, toxicity of to
cabbage looper, 991
Chrysopa spp., 70
convergent lady beetle, 70
corn earworm, 22
Collops vittatus, 70
Geocoris spp., 70
German cockroach, 1008
house fly, 206, 233, 709, 1005
milkweed bug, 1007
mouse, 206
Nabis ferus, 70
oriental fruit moth, 469
Orius sp., 70
rats, 233

salt-marsh caterpillar, 1022
Sex lure, natural female, for detection of gypsy moth, 751
Sheep
effect of dieldrin on, 826
malathion residues in tissues of, 972
Shell SD-3562
effects of on germination of wheat, 575
effect of on beetle, 1035
synergistic action of, 887
systemic action in cortical tissues of elm, 1034
systemic effectiveness of against bed bug, 6
systemic effectiveness of against Gulf Coast tick, 6
Shell SD-3562, against
boll weevil, 747
bollworm, 747
cotton leafworm, 747
desert spider mite, 748
Shell SD-3562, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Shell SD-3562, for control of
boll weevil, 1052
cotton aphid, 747
cotton fleahopper, 1052
introduced pine sawfly, 648
spider mites, 1053
thrips, 1051
Shell, SD-3562, toxicity of to
house fly, 6, 889
white rat, 6
Shell SD-4402
long term effectiveness against eye gnat, 785
loss of from soil surface, 650
toxicity of to corn earworm, 22
Shell SD-4402, against
boll weevil, 747
bollworm, 747
cotton leafworm, 747
eye gnat, 367
Shell SD-4402, for control of
army cutworms, 319
bean leaf skeletonizer, 566
boll weevil, 1052
cotton fleahopper, 1052
European pine shoot moth, 352
green peach aphid, 1099
lesser cornstalk borer, 664
spider mites, 1053
thrips, 1051
Shell SD-5539
against boll weevil, 747
against bollworm, 747
for control of tarnished plant bug, 690
SG 67
effect of on cat flea, 906
effect of on oriental rat flea, 906
effect of on short-nosed cattle louse, 906
for control of snake mite, 903
Silesis musculus, 1018
Silica aerogel, for control of tropical rat mite, 475
Silverfish, malathion for reducing dried-fruit packing house infestations of, 970
Simuliidae, effect of DDT on, 609
Sinea diadema, 69
Siphona irritans, 435, 477
Sitona hispidula, 537, 865
Sitophilus
granarius, 341, 375, 537
oryza, 215, 238, 288, 341, 372, 456, 537, 745, 1071
Sitotroga cerealella, 238, 536, 573, 973
Sitticus sp., 813
Smilia camelus, 173
Sminthuridae, abundance of in dieldrin-treated areas, 823
Soapstone, toxicity of to corn earworm, 22
Sodium arsenate, in baits for control of imported fire ant, 190
Sodium arsenite, in baits for control of imported fire ant, 190
Sodium cyanide, effect of on transamination in cockroach, 167
Sodium fluoride, in baits for control of imported fire ant, 190
Sodium fluoroacetate, in baits for control of imported fire ant, 190
Sodium fluosilicate
effectiveness of on salt-marsh caterpillar, 78
for control of desert grasshopper, 724
Sodium orthoarsenate, effect of on transamination in cockroach, 167
Soil, influence of on beet webworm, 585
Soil injection, as means of applying systemic acaricides to fruit trees, 144
Soil pH, effect of on grub populations, 860
Soil treatment, in lieu of spraying for control of plum curculio, 439
Soil type, effect of on germination and growth of wheat, 771
Solenopsis
saevissima, 1137
saevissima v. *richteri*, 188, 646
Solierella sp., 958
Solvents
Ansco, influence of on efficiency of granulated insecticides, 943
AR Series, influence of on efficiency of granulated insecticides, 940
Espesol, influence of on efficiency of granulated insecticides, 941
Panasol, influence of on efficiency of granulated insecticides, 945
Penola, influence of on efficiency of granulated insecticides, 946
Shell Oil Company, influence of on efficiency of granulated insecticides, 943
Standard Oil of California, influence of on efficiency of granulated insecticides, 941
Sorghum grass, effect of hydrogen phosphide on seed germination of, 1
Soybean
effect of absorbents on, 491
effects of soil insecticides on, 491
observations of insects on, 131
Sevin residues on, 481
Sparganothis sulfureana, 527
Spermatozoa, honey bee, effect of drying and various diluents on, 266
Spiders, effect of insecticides on, 358
Spinach, residues of nicotine on, 64
Spodoptera exigua, 616, 729
Spores, Nosema, longevity of refrigerated, 1132
Spraying, aerial
against spruce budworm, 631
for control of maple leaf cutter, 957
Stalk shredding, for control of pink bollworm, 437
Stannic chloride, against house fly, 447
Stannous octanoate, against house fly, 447
Stannous oleate, against house fly, 447
Starch, growth and survival of mosquito larvae on, 169
Starvation tests, with larvae of cinnabar moth, 472
Stauffer R-2371, effectiveness of against cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Stauffer R-2448, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Stenonema, 609
Stephanitis
globulifera, 406
pyrioides, 406, 839
rhododendri, 406
Stethorus, 982
Stimulant, as standard term for insect reaction to chemicals, 134
Stomoxys calcitrans, 301, 408, 432, 692, 850, 1038, 1125
Stored-grain insects, control of, 536
Strobane
against boll weevil, 747
against bollworm, 747
residues of in mint oil, 531
toxicity of to oriental fruit moth, 470
Strobane, for control of
boll weevil, 551, 1051
cotton fleahoppers, 1051
European pine shoot moth, 366
mint insects, 529
Nantucket pine moth, 366
plum curculio, 335
spider mites, 1053
thrips, 1051
Sugar, growth and survival of mosquito larvae on, 169
Sugar, plant
effect of DDT on, 492
effect of Hydrodarco B on, 492
Sugar beet, effect of virus yellows on yield and quality of, 162
Sulfate of 4,5-diaminouracil, effect of on house fly development, 803
Sulfoxide, as synergist for barthrin in sprays against house fly, 282
Sulfur
for control of citrus rust mites, 396
for control of grape lead folder, 533
for control of lygus bugs, 398
influence of on growth of apple, 717
Sulfuryl fluoride, fumigation of buildings with, 508
Sulphenone, for control of a spider mite complex, 522
Sulphur dust, for control of a spider mite complex, 522
Supella supellectilium, 904
Survival
effect of low temperature on in alfalfa weevil, 570
of wheat stem sawfly, relationship between culm solidness and, 246
Swine, effect of dieldrin on, 826
Synanthedon pictipes, 321
Synapse, insect, action of DDT at, 171
Syneura cocciphila, 1057
Syrphidae, as predators of spotted alfalfa aphid, 92
Syrphids, toxicity of insecticides to, 69
Syrphus nitripennis, 659
Systena
blanda, 527, 823
elongata, 132
frontalis, 527
Systox, toxicity of to house fly, 742
Sytam, for control of holly scale, 406

Tabatrex
for control of face fly, 450
repellency of to horn fly, 435

repellency of to stable fly, 434
Tabutrex, toxicity of to corn earworm, 22
Tartar emetic, in baits for control of imported fire ant, 190
TDE
effect of on hornworms, 976
solubilization of by lipoprotein, 34
toxicity of to corn earworm, 22
toxicity of to house fly, 38
TDE, for control of
blackheaded fireworm of cranberry, 1097
cranberry fruitworm, 1118
green peach aphid, 1099
mint insect, 528
mites on apple, 735
TDE, aliphatic deuterium analog of degradation of when applied to
house fly, 40
synthesis of, 35
toxicity of to house fly, 35
Techniques (*see Methods*)
Tedion
against two-spotted spider mite, 841
effect of on demeton-resistant citrus red mites, 388
effect of on ovex-resistant citrus red mites, 388
toxicity of to oriental fruit moth, 469
Tedion, for control of
cotton mite, 204
European red mite, 685
mites on apple, 735
Oligonychus pratensis, 1061
spider mite complex, 522
two-spotted spider mite, 967
Temperature
low, effect of on eggs of Angoumois grain moth, 973
low, effect of on eggs of confused flour beetle, 973
low, effect of on eggs of Indian-meal moth, 973
low, effect of on survival of alfalfa weevil, 570
low, response of quiescent khapra beetle larvae to, 698
relation to moisture content in effectiveness of fumigants in stored grain, 1071
storage, influence on effectiveness of malathion, 341
varying, effect of on diapause in pink bollworms, 794
Tenebrio
molitor, 382, 537
obscurus, 537
Tenebroides mauritanicus, 239, 537
TEPP, for control of
European pine shoot moth, 366
Nantucket pine moth, 366
Tetra-*n*-butyltin, against house fly, 447
Tetra-*n*-lauryltin, against house fly, 447
Tetra-*n*-octyltin, against house fly, 447
Tetraethyltin, against house fly, 447
Tetragnatha sp., 813
Tetraisopropyltin, against house fly, 447
Tetralopha robustella, 587
Tetramethylene methanesulfonate, effect of on house fly development, 803
Tetramethyltin, against house fly, 447
Tetranychus, 131
atlanticus, 113, 204
bimaculatus, 389
cinnabarinus, 105, 725
desertorum, 54, 747, 1051
mcdanieli, 522
pacificus, 204, 626, 982
telarius, 80, 127, 144, 204, 224, 487, 491, 522, 527, 661, 735, 841, 967
Tetraphenyltin, against house fly, 447
Tetravinyltin, against house fly, 447
Tettigoniidae, abundance of in dieldrin-treated areas, 823
Thallium sulfate, in baits for control of imported fire ant, 190
Thamnosphecia pyri, 276
Thanite, toxicity of to
Chrysops, 263
house fly, 741
Theridiidae, in apple orchards, 811
Theridion
albidum, 812
differens, 812
murarium, 811
rabuni, 812
Therioaphis
maculata, 71, 89, 93, 142, 146, 220, 234, 257, 655, 667, 714, 1094
trifolii, 113
Thimet (*see Phorate*)
Thioammelin, effect of on house fly development, 803
Thiodan
against boll weevil, 711
against green peach aphid, 624
as soil treatment against eye gnat, 1102
effect of on lesser cornstalk borer, 665
for bee poisoning in red clover, 1013
persistance of residues on alfalfa, 960
Thiodan, for control of
army cutworms, 319
bean leaf skeletonizer, 566
boll weevil, 229
clover aphid, 1013
corn earworm, 819
cranberry fruitworm, 1116
Douglas-fir beetle, 949
eye gnats, 362
green peach aphid, 403, 725, 871, 963, 1099
Hyperodes humilis, 254
lesser peach tree borer, 321
lygus bugs, 74, 767
pea aphid, 819
plum curculio, 439
potato tuberworm, 868
spider mites, 726
three-lined potato beetles, 480
Thiodan, toxicity of to
cabbage looper, 991
corn earworm, 22
house fly, 741
oriental fruit moth, 469
Thiourea, effect of on fertility of house fly, 804
Thiram, effect of on germination of wheat, 575
Thomisidae, in apple orchards, 811
Thrips
nigropilosus, 527
tabaci, 131, 527
Thuricide, in baits for control of imported fire ant, 190
Thyanta custator, 689
Thymoxyacetic acid, effect of on house fly development, 804
Thymine, effect of on house fly development, 803
Thysanus, 907
Timberlakia europaea, 694
Tipula mingwe, 527
Tmarus sp., 813
Tobacco, suitability of various for growth of cigarette beetle, 381
Tomato, reaction of *Drosophila* to varieties of, 337
Torymus druparum, 166
Toumeyella numismaticum, 325
Toxaphene
effect of on transamination in cockroach, 167
seasonal tolerance of boll weevil to, 41
susceptibility of boll weevil to, 311
Toxaphene, against
boll weevil, 711, 747
bollworm, 747
cotton leafworm, 749
European corn borer, 625
eye gnat, 367
stable fly, 303
Toxaphene, effectiveness of against
boll weevil, 53
corn earworm, 54
cotton fleahopper, 53
desert spider mite, 54
salt-marsh caterpillar, 78
Toxaphene, effect of on
boll weevil, 294
Ladino clover, 1134
lesser cornstalk borer, 665
Sacadodes, 294
Toxaphene, for control of
bean leaf skeletonizer, 566
boll weevil, 229, 1051
corn earworm, 397
cotton fleahopper, 1051
cotton mite, 204
desert grasshopper, 724
eye gnats, 365
face fly, 450
fuller rose beetle, 164
European corn borer, 590
European pine shoot moth, 366
green peach aphid, 403
Hyperodes humilis, 253
imported fire ant, 190
lone star tick, 953
lygus bugs, 397
mint insects, 527
morning-glory leaf miner, 1119
Nantucket pine moth, 366
spider mites, 1053
sugar-beet wireworm, 177
thrips, 227, 1051
Toxaphene, residues of in/on
forage under pecan, 719
hogs, 322
mint oil, 527
potato, 833
radish, 833
sugar beet, 832
table beet, 832
Toxaphene, toxicity of to
boll weevil, 534, 551
cabbage looper, 99
Chrysopa spp., 69
Chrysops, 263
convergent lady beetle, 69
corn earworm, 22
Geocoris spp., 69
Nabis ferus, 69
Orius sp., 69
Sinea diadema, 69
spotted alfalfa aphid, 69
syrphids, 69
Toxoptera graminum, 199, 258, 278, 454, 473, 604, 798
Trachelas tranquillus, 813
Transamination, effect of insecticides on in cockroach, 166
Transport, aerial
of greenbug, 604
of harlequin bug, 604
Trap
for deciduous tree fruit flies, 967
light, for mosquitoes, 876
Graham, in gypsy moth detection, 751

to recover dead and abnormal honey bees, 782
Triacetin, effect of on house fly development, 804
Trialeurodes abutilones, 104
Triatoma protracta 904
Tribolium, 970
castaneum, 170, 238, 496, 537, 974
confusum, 341, 382, 537, 691, 762, 973, 974, 1071
Tri-*n*-butyltin chloride, against house fly, 447
Tri-*n*-butyltin-dibutoxy-dithiophosphate, against house fly, 447
Tri-*n*-butyltin octanoate, against house fly, 447
Tricalcium arsenate, effect of on boll weevil, 845
Trichogramma minutum, 304
Trichoplusia ni, 47, 77, 104, 242, 389, 590, 991
Trichoptera, attracted to black light, 173
N, *N'*, *N''*-Triethylenephosphorothioic triamide, effect of on
fertility of house fly, 804
house fly development, 804
Triethyltin acetate, against house fly, 447
Triethyltin benzyl mercaptide, against house fly, 447
Triethyltin ethyl mercapto acetate, against house fly, 447
Triethyltin hydroxide against house fly, 447
Triethyltin phenyl mercaptide, against house fly, 447
Trigonotylus ruficornis, 527
Triisopropyltin acetate, against house fly, 447
Triisopropyltin undecylenate, against house fly, 447
Trimedlure, added common name of insecticide, 677
Trimerotropis pallidipennis pallidipennis, 721
Trimethyltin bromide, against house fly, 447
Trioxys utilis, 69, 655, 1094
Triphenyltin chloride, against house fly, 447
Tris-(tri-*n*-butyltin) perthiophosphate, against house fly, 447
Trithion
against desert spider mite, 748
as soil treatment against eye gnat, 1102
duration of toxicity of to fuller rose beetle, 165
effect of on survival of serpentine-leaf-miner parasite, 873
for bee poisoning in red clover, 1013
honey bee mortality from, 1014
persistence of in soil, 1080
systemic effectiveness of to bed bug, 6
systemic effectiveness of to Gulf Coast tick, 6
Trithion, for control of
bean leaf skeletonizer, 567
boll weevil, 230
clover aphid, 1012
codling moth, 1083
corn earworm, 401
cotton aphid, 747
cotton leaf perforator, 76
cranberry fruitworm, 1116
European pine shoot moth, 352, 365
grape mealybug, 356
Hyperodes humilis, 256
lygus bugs, 401, 514, 767
Nantucket pine moth, 365
Oligonychus pratensis, 1061
plum curculio, 335
poultry lice and mites, 161
San Jose scale, 1085
sap beetles, 175
serpentine leaf miner, 873
spider mite complex, 522
spider mites, 725
tarnished plant bug, 690
variegated cutworm, 74
woolly apple aphid, 217
Trithion, toxicity of to
corn earworm, 22
house fly, 6
oriental fruit moth, 470
white rat, 6
Tritneplis herocompae, 305
Triton X-100, for control of *Hyperodes humilis*, 256
Triton X-151, in phytotoxic formulations, 203
Triton X-171, in phytotoxic formulations, 203
Trivinyltin iodide, against house fly, 447
Trogoderma
granarium, 698
inclusum, 537
Tropic acid, as synergist in house fly sprays, 300
Trybliographa rapae, 65, 933
Tyrophagus, 904
Tyrosine, effect of on house fly development, 804
Typhaea stercorea, 536
Tyria jacobaeae, 472

UCC-8305
systemic effectiveness of against bed bug, 6
systemic effectiveness of against Gulf Coast tick, 6
toxicity of to house fly, 6
toxicity of to white rat, 6
Underfeeding, for prolongation of life of mosquito larvae, 169
Uracil, effect of on house fly development, 804
Uracil-4-acetic acid, effect of on house fly development, 804
Urea, in control of apple insects, 685

VC-13, systemic and topical activity of, 692
Van De Graaff electron accelerator, use of in dosimetry studies, 496
Vapam, fumigation of soil with, 1080
Vaporizer, fumigant, for laboratory use, 562
Ventilation unit, for aerating insectary sleeve cages, 959
Vinyl di-*n*-dibutyltin chloride, against house fly, 447
Virginia-Carolina 1-13, effectiveness of against
cattle grubs, 1126
lone star tick, 1126
screw-worms, 1126
stable fly, 1126
Virus yellows, effect of on yield and quality of sugar beet, 162

Water, effect of on transamination in cockroach, 167
Whale meat, for mass-production of screw-worms, 1114
Wheat
chloroplast number in Hessian-fly infested leaves, 560
chloroplast number in leaves treated with maleic hydrozide, 560
development of granary weevils and storage fungi in, 375
distribution and sorption of liquid fumigants applied to by recirculation, 259
effect of hydrogen phosphide on seed germination of, 1
effect of soil type and moisture on germination and growth of, 771
effect of systemic insecticides on germination of, 575
estimating rice weevil populations in, 215
fumigation of for control of rice weevil, 288
measurements of greenbug damage to, 798
relation between silica in and resistance to Hessian fly, 995
stored, fumigation of, 537, 1071
Wheat germ
effect of on house fly, 325
medium, for rearing pink boll worm, 759

Xenopsylla cheopis, 167, 170, 247, 904
Xylene, in phytotoxic formulations, 203
Xyletinus peltatus, 329
Xylocopa virginica, 679
Xylocoris galactinus, 537
Xysticus sp., 813

Yellow dwarf virus, and greenbug dispersal, 473
Yield
beans, effect on of various insecticides for control of lygus bugs and corn earworm, 397
clover, relation of to bee counts, 549
corn, effect of European corn borer on, 517
corn, effect of sugarcane borer infestation on, 935
corn, loss of associated with corn leaf aphid, 924
cotton, effect of insecticides on, 291
mushroom, from beds treated with lindane, 558
red clover, in clover aphid control plots, 1013
sorghum, effect of phorate applied to seed on, 154
sugar beet, effect of virus yellows on, 162
vegetable, effect of Sevin on, 388
wheat, influence of soil type and moisture on, 773

Zabrotes subfasciatus, 239
Zineb, for control of
citrus rust mite, 395
mint insects, 528
mites on apple, 735
Zygogramma suturalis casta, 527